



May 24, 2024

Serapure Preparation and Testing

 Forked from [DNA EXTRACTION Protocol Template](#)

This protocol is a draft, published without a DOI.

Andreas Novotny¹

¹University of British Columbia

Better Biomolecular Ocea...



Andreas Novotny

University of British Columbia, Hakai Institute

OPEN  ACCESS



Protocol Citation: Andreas Novotny 2024. Serapure Preparation and Testing. [protocols.io](https://protocols.io/view/serapure-preparation-and-testing-dc5x2y7n) <https://protocols.io/view/serapure-preparation-and-testing-dc5x2y7n>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development

We are still developing and optimizing this protocol

Created: November 07, 2023

Last Modified: May 24, 2024

Protocol Integer ID: 99223

Disclaimer

Draft!

Abstract

A protocol template created through the BeBOP project for DNA Extraction.

Guidelines

MIOP: Minimum Information about an Omics Protocol

MIOP Term	Value
analyses	
audience	
broad-scale environmental context	
creator	
environmental medium	
geographic location	
hasVersion	
issued	
language	
license	
local environmental context	
materials required	
maturity level	
methodology category	
personnel required	
project	
publisher	
purpose	
skills required	
target	
time required	

See <https://github.com/BeBOP-OBON/miop/blob/main/model/schema/terms.yaml> for list and definitions.

AUTHORS

PREPARED BY All authors known to have contributed to the preparation of this protocol, including those who filled in the template.	AFFILIATION	ORCID (visit https://orcid.org/ to register)	DAT
Content Cell	Content Cell	Content Cell	yyyy-mm-dd
Content Cell	Content Cell	Content Cell	yyyy-mm-dd

This protocol is derived from the referenced protocol created by Nadin Rohland. Please do not cite this protocol as being originally produced by me.

RELATED PROTOCOLS

PROTOCOL NAME AND LINK	ISSUER / AUTHOR	RELEASE / ACCESS DATE
Content Cell	Content Cell	yyyy-mm-dd
Content Cell	Content Cell	yyyy-mm-dd

This is a list of other protocols which should be known to users of this protocol. Please include the link to each related protocol.

Suggested Reading

Rohland N, Reich D. **Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture.** Genome Research. Early Online Access. Doi: [10.1101/gr.128124.111](#)

DeAngeliss MM, Wang DG, Hawkins TL: **Solid-phase reversible immobilization for the isolation of PCR products.** *Nucleic Acids Res* 1995, **23**:4742–4743.

Fisher S, et al.: **A scalable, fully automated process for construction of sequence-ready human exome targeted capture libraries.** *Genome Biol* 2011, **12**:R1.

Lundin S, Stranneheim H, Pettersson E, Klevebring D, Lundeberg J: **Increased throughput by parallelization of library preparation for massive sequencing.** *PLoS One* 2010, **5**:e10029.

ACRONYMS AND ABBREVIATIONS

ACRONYM / ABBREVIATION	DEFINITION
Content Cell	Content Cell

GLOSSARY

SPECIALISED TERM	DEFINITION
Content Cell	Content Cell
Content Cell	Content Cell

BACKGROUND

This document describes the required protocol to conduct insert name of the method/protocol.

Summary

The goal here is to create a substitute for AMPure XP that is of equal effectiveness in comparison to the commercial product but far more cost-effective (\$19/mL versus \$0.46/mL).

Insert a short description of the background for the method/protocol (e.g. why and for which purpose do you perform water sampling). Please provide a brief summary of your method including, as appropriate, a brief description of what techniques your best practice is about, which ocean environments or regions it targets, the primary sensors covered, what type of data/measurements/observing platform it covers, limits to its applicability.

Method description and rationale

Insert a short description of the functioning principal of the methodology used in the protocol (i.e. how does the method work?). Please note that this is different from the step-by-step description of the protocol procedure. Insert a short statement explaining why the specific methodology used in the protocol has been selected (e.g. it is highly reproducible, highly accurate, procedures are easy to execute etc....).

Spatial coverage and environment(s) of relevance

If applicable, please specify the region where the protocol is applied. For regional term guidance see here. If applicable, please indicate here the environment(s) of relevance for the protocol, e.g. Abyssal plain. Select from the ENVO terminology.

Personnel Required

Insert the number of technicians, data managers, and scientists required for the good execution of the procedure

Safety

Identify hazards associated with the procedure and specify protective equipment and safety training required to safely execute the procedure

Training requirements


Specify technical training required for the good execution of the procedure.

Time needed to execute the procedure

Specify how much time is necessary to execute the procedure.

Materials


DESCRIPTION e.g. filter	PRODUCT NAME AND MODEL Provide the official name of the product	MANUFACTURER Provide the name of the manufacturer of the product.	QUANTITY Provide quantity
Durable equipment			
Content Cell	Content Cell	Content Cell	Content Cell
Content Cell	Content Cell	Content Cell	Content Cell
Consumable equipment			
Content Cell	Content Cell	Content Cell	Content Cell
Content Cell	Content Cell	Content Cell	Content Cell
Chemicals			
Content Cell	Content Cell	Content Cell	Content Cell
Content Cell	Content Cell	Content Cell	Content Cell

 Sera-Mag Speedbeads, magnetic carboxylate modified particles (Cytiva 65152105050250) **Thermo Fisher Scientific Catalog #09-981-123**

Note

These are the same beads used by Beckman, per Orapure product sheet and <http://bit.ly/vmiDzU>

- PEG-8000 (Amresco 0159)
- 0.5 M EDTA, pH 8.0 (Amresco E177)
- 1.0 M Tris, pH 8.0 (Amresco E199)
- Tween 20 (Amresco 0777)
- 5 M NaCL

 GeneRuler Ultra Low Range DNA Ladder **Thermo Fisher Catalog #SM1211**

 6-Tube Magnetic Separation Rack **New England Biolabs Catalog #S1506S**

Optional

- Agencourt SPRIPlate Super Magnet Plate (Beckman Coulter A32782)

Before start

Read background information, MIOP and BePOP-OBON information under the "Guidelines" tab.















STANDARD OPERATING PROCEDURE



- 1 In the following SOP, please use the exact names of equipment as noted in the table above.

Provide a step-by-step description of the protocol. The identification of difficult steps in the protocol and the provision of recommendations for the execution of those steps are encouraged.

STEPS


- 2 In a 50 mL conical using sterile stock solutions, prepare TE ([M] 10 millimolar (mM) TrisNHCl, [M] 1 millimolar (mM) EDTA =  500 µL [M] 1 Molarity (M) Tris pH8 +  100 µL [M] 0.5 Molarity (M) EDTA, fill conical to 50 mL mark with dH2O).
- 3 Mix SeraNmag SpeedBeads and transfer  1 mL to a 1.5 mL microtube.
- 4 Place SpeedBeads on magnet stand until beads are drawn to magnet.
- 5 Remove supernatant with P200 or P1000 pipetter.
- 6 Add  1 mL TE to beads, remove from magnet, mix, return to magnet.
- 7 Remove supernatant with P200 or P1000 pipetter.
- 8 Add  1 mL TE to beads, remove from magnet, mix, return to magnet.
- 9 Remove supernatant with P200 or P1000 pipetter.
- 10 Add  1 mL TE to beads and remove from magnet. Fully resuspend and set microtube in rack (i.e. not on magnet stand).
- 11 Add  9 g PEGN8000 to a new 50 mL, sterile conical.
- 12 Add  10 mL [M] 5 Molarity (M) NaCl (or  2.92 g) to conical.
- 13 Add  500 µL [M] 1 Molarity (M) TrisNHCL to conical.
- 14 Add  100 µL [M] 0.5 Molarity (M) EDTA to conical.
- 15 Fill conical to ~ 49 mL using sterile dH2O. You can do this by eye, just go slowly.
- 16 Mix conical for about 3N5 minutes until PEG goes into solution (solution, upon sitting, should be clear).
- 17 Add  27.5 µL Tween 20 to conical and mix gently.






- 18 Mix  1 mL SpeedBead + TE solution and transfer to 50 mL conical.
- 19 Fill conical to 50 mL mark with dH2O (if not already there) and gently mix 50 mL conical until brown.
- 20 Test against AMPure XP using aliquots of ladder (Fermentas GeneRuler). I recommend the 50 bp ladder in place of the ultraNlow range ladder.
- 21 Wrap in tinfoil (or place in dark container) and store at  4 °C .

22

Note

You may also wish to prep an extra  50 mL of PEG solution that lacks SeraNmag SpeedBeads so that you can use it in a bead-inclusive library preparation protocol, derived from Fisher (2011).

In that case, just:

1. Add  10 g PEGN8000 to a new 50 mL, sterile conical.
2. Add  25 mL [M] 5 Molarity (M) NaCL (or  7.3 g) to conical.
3. Fill conical to ~ 49 mL using sterile dH2O. You can do this by eye, just go slowly.
4. Mix conical for about 3-5 minutes until PEG goes into solution (solution, upon sitting, should be clear).







- 23 Test monthly.

TESTING

24









Note

You should test the Serapure mixture to ensure that it is working as expected. You can do this using DNA ladder (Fermentas GeneRuler – NEB ladders may cause problems).

- 25 Prep fresh aliquots of [M] 70 Mass / % volume EtOH.
- 26 Mix  2 µL GeneRuler with  18 µL dH2O.
- 27 Add  20 µL GeneRuler mixture to a volume of Serapure and/or AMPure (the specific volume depends on whether you are trying exclude small fragments or not; see the figure on the next page).
- 28 Incubate mixture  00:05:00 at  Room temperature .
- 29 Place on magnet stand.
- 30 Remove supernatant.
- 31 Add  500 µL [M] 70 Mass / % volume EtOH.


5m



- 32  0 μL Incubate on stand for  00:01:00 . 1m
- 33 Remove supernatant.
- 34 Add  500 μL [M] 70 Mass / % volume EtOH.
- 35 Incubate on stand for  00:01:00 . 1m
- 36 Remove supernatant.
- 37 Place beads on  37 $^{\circ}\text{C}$ heat block for 3-4 min. until dry.
- 38 Rehydrate with  20 μL dH₂O.
- 39 Place on magnet stand.
- 40 Transfer supernatant to new tube.
- 41 Mix supernatant with  1 μL loading dye.
- 42 Electrophorese in 1.5 % agarose for  01:00:00 at 100 V. 1h

QUALITY CONTROL

- 43 The following image compares the results of “purifying” a mix of 2 μL Fermentas Ultra Low Range Ladder + 18 μL dH₂O using several different amounts of AMPure or SeraPure solution to DNA solution. AMPure is on the left, “SeraPure” is on the right. After preparing 20 μL of ladder + water mix, we combined that with the volumes of AMPure or SeraPure listed below and then purified using the standard protocol:



As you can see, the volume of AMPure or SeraPure controls the size of fragments recovered. More specifically, it is the ratio of PEG solution used to the volume of the DNA in solution which makes the difference, not the count of beads in solution (provided they are above the minimum level). This is what makes it possible to do “doubleNSPRI” size selection.



BASIC TROUBLESHOOTING GUIDE

- 44 Identify known issues associated with the procedure, if any.

Provide troubleshooting guidelines when available.

APPENDIX A: DATASHEETS

- 45 Link templates (e.g. preformatted spreadsheets) used to record measurements and report on the quality of the data as well as any documents such as manufacturer specifications, images, etc that support this protocol. Please include a short note describing the document's relevance.

Protocol references

Insert all references cited in the document.

Please insert full DOI address when available, e.g. <http://doi.dx.org/10.1007/s11258-014-0404-1>