




Upload image

May 14, 2020

Western blot analysis and immunoprecipitation assay

Marzia Ognibene¹¹Laboratorio Cellule Staminali Post Natali e Terapie Cellulari, IRCCS Istituto Gaslini, Genova, Italy**1** Works for me dx.doi.org/10.17504/protocols.io.bgecjtaw Marzia Ognibene

- 1 Cells are washed in cold PBS and lysed on ice in a buffer containing 1.6 mM NaH₂PO₄, 8.6 mM Na₂HPO₄, 1% Triton X-100, 0.1% SDS, 0.1% NaCl, 0.5% NaDoc, 2 mM AEBSF, 20 mg/mL each of aprotinin and leupeptin.
- 2 Cell lysates are centrifuged in a microfuge at maximum speed for 10 min to eliminate debris.
- 3 Protein content is evaluated by bicinchoninic acid assay (BCA Protein assay kit by Thermo Fisher) in a spectrophotometer.
- 4 Lysates (100 µg each) are subjected to SDS-PAGE electrophoresis and transferred to PVDF membrane.
- 5 For the IMMUNOPRECIPITATION ASSAY, cells are lysed in a buffer containing 20 mM Tris-HCl, 1% Triton X-100, 10% Glycerol, 2 mM AEBSF, 20 mg/mL each of aprotinin and leupeptin.
- 6 Lysates (1000 µg) are incubated overnight at 4°C with the desired antibody under constant rotation.
- 7 Gammabind G-Sepharose (40 µl) is added to each sample and let rotate for 1 hour and 30 min.
- 8 Samples are washed three times with the lysis buffer in a microfuge at 4300 g for 1 min.
- 9 After eliminating the last supernatant, samples are eluted in 40 µl of 1x Laemmli's buffer, subjected to SDS-PAGE and transferred to PVDF membrane.
- 10 PVDF membrane is incubated in blocking buffer (4% nonfat milk in T-TBS -Tris Buffer Solution 1x with 0,05%Tween) for 2 hr at room temperature.
- 11 Membrane is incubated in primary antibody diluted in 0,5%BSA in T-TBS, overnight at 4°C.

- 12 Membrane is washed in T-TBS, 5 times, 5 min each.
- 13 Membrane is incubated in HRP-conjugated secondary antibody diluted in 0,5%BSA in T-TBS, 2 hr, at room temperature.
- 14 Membrane is washed in T-TBS, 5 times, 5 min each.
- 15 Proteins are visualized by enhanced chemiluminescence (ECL) detection, through the Bio-Rad ChemiDoc instrument. Signal intensity was measured by densitometry using the Bio-Rad Image Lab 6.0 software.