

Jul 09, 2024

© DEVELOPMENT OF A QUESTIONNAIRE OF COMMITMENT TO NEW PHYSICAL INTERVENTIONS FOR ATHLETES: DELPHI RESEARCH

DOI

dx.doi.org/10.17504/protocols.io.14egn6w5ql5d/v1

Rybakova Polina Denisovna¹, Miroshnikov Alexander Borisovich¹, Shevtsov Alexey Victorovich¹ Russian University of Sports "GTSOLIFK"



Polina Rybakova

Russian university of sports "GTSOLIFK"





DOI: dx.doi.org/10.17504/protocols.io.14egn6w5ql5d/v1

Protocol Citation: Rybakova Polina Denisovna, Miroshnikov Alexander Borisovich, Shevtsov Alexey Victorovich 2024. DEVELOPMENT OF A QUESTIONNAIRE OF COMMITMENT TO NEW PHYSICAL INTERVENTIONS FOR ATHLETES: DELPHI RESEARCH. **protocols.io** https://dx.doi.org/10.17504/protocols.io.14egn6w5ql5d/v1

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's
working

Created: July 09, 2024

Last Modified: July 09, 2024

Protocol Integer ID: 103075

Keywords: questionnaire, Delphi method, physical activity, physical performance, sport, athlete



Abstract

In the context of exercise, commitment is the correct level of a person's performance of a recommended training regimen (for example, the number of training days per week), including the level at which a person follows the prescribed dosage of exercise (for example, determined by heart rate, perceived load level, or percentage of one repeated maximum). There is no questionnaire to assess exercise adherence for athletes. The consensus of experts will be used to create the questionnaire. to develop a questionnaire to assess adherence to a new training effect for athletes using the Delphi method.

Attachments



Guidelines

The study used the Delphi method to display the current consensus among experts. The CREDES (Conducting and REporting Delphi Studies) checklist was used to ensure compliance with reporting rules and the ACCORD (ACcurate Consensus Reporting Document) reporting guideline is dedicated to and applicable to all consensus methods used in biomedical research. The Delphi method is recommended as a reliable way to achieve consensus among experts. Delphi was planned to consist of three rounds to allow for iteration, with the ability to include additional rounds as needed.

Before start

Hypothesis: Valid and reliable assessment of exercise adherence results will allow for informed conclusions about the effects of exercise.

Round 1. According to ACCORD (items M6 and M7), we conducted a systematic literature search to create the first questionnaire on adherence to new exercise interventions for athletes [5]. Four experts reached a consensus to include 39 questions in the first questionnaire. The expert council discussed the resulting questionnaire after the first round and jointly developed a protocol for the second and third rounds. The first round protocol was registered in the OSF international database: https://doi.org/10.17605/OSF.IO/SA2E9.

Round 2. The protocol for the second and third rounds was compiled before the start of the search and did not change either during or after its completion, according to ACCORD and PRISMA-P (Preferred Reporting Items for Systematic Review and Meta Analysis Protocols (PRISMA-P))



Materials

Experts' choice. Participants were recruited using purposive sampling, which selected a panel of experts based on their knowledge and experience in sport, scientific interest and communication skills. The process of selecting experts began with one of the authors (A.V. Shevtsov) sending an email invitation to experts who met the selection criteria. This email contained information about the research project, as well as informed consent forms and a 39-question questionnaire. Experts were invited to contact the author by email or telephone to discuss the project. Participants were also invited to forward information about the project to interested sports professionals who they believed would meet the selection criteria. Interested experts then confirmed their participation via email. Ethics committee approval was not required for this study. All participants provided written informed consent prior to inclusion in the study. As described in ACCORD, there are no generally accepted standards for the number of experts recruited in Delphi studies, although 20 to 30 is common. Thus, the target panel size (about 20 participants) was determined by the desired representation and ensuring an acceptable response rate (10, assuming a 50% participation rate) in the event of nonparticipation or partial completion of the survey. Panelists who agreed to participate received an information pack containing an introductory letter, a plain language summary, an informed consent statement, links to the published protocol and systematic review, and a 39question questionnaire.

We invited 30 people to participate in the Delphi process.

Before start

Hypothesis: Valid and reliable assessment of exercise adherence results will allow for informed conclusions about the effects of exercise.

Round 1. According to ACCORD (items M6 and M7), we conducted a systematic literature search to create the first questionnaire on adherence to new exercise interventions for athletes [5]. Four experts reached a consensus to include 39 questions in the first questionnaire. The expert council discussed the resulting questionnaire after the first round and jointly developed a protocol for the second and third rounds. The first round protocol was registered in the OSF international database: https://doi.org/10.17605/OSF.IO/SA2E9.

Round 2. The protocol for the second and third rounds was compiled before the start of the search and did not change either during or after its completion, according to ACCORD and PRISMA-P (Preferred Reporting Items for Systematic Review and Meta Analysis Protocols (PRISMA-P))





Protocol references

Slade SC, Dionne CE, Underwood M, Buchbinder R. Consensus on Exercise Reporting Template (CERT): Explanation and Elaboration Statement. Br J Sports Med. 2016 Dec;50(23):1428-1437. doi: 10.1136/bjsports-2016-096651.

Bailey DL, Bishop A, McCray G, Foster NE, Holden MA. A new measure of exercise adherence: the ATEMPT (Adherence To Exercise for Musculoskeletal Pain Tool). Br J Sports Med. 2024 Jan 3;58(2):73-80. doi: 10.1136/bjsports-2022-106347.

Taylor L, Wiebusch M, Bisset LM, Coombes BK. Adherence to exercise in lateral elbow tendinopathy, a scoping review. Musculoskelet Sci Pract. 2024 May 21;72:102978. doi: 10.1016/j.msksp.2024.102978.

Yang Y, Gao Y, An R, Wan Q. Barriers and facilitators to exercise adherence in community-dwelling older adults: A mixed-methods systematic review using the COM-B model and Theoretical Domains Framework. Int J Nurs Stud. 2024 May 21;157:104808. doi: 10.1016/j.ijnurstu.2024.104808.

Шевцов, А. В. Обзор литературы, предметного поля для разработки опросника приверженности к физическим упражнениям спортсменов: первый раунд метода Delphi/A. В. Шевцов, А. Б. Мирошников, В. В. Большаков, П. Д. Рыбакова // Спортивно-педагогическое образование. – 2024. – № 2. – С. 50–58.

Levites Strekalova YA, Nelson JD, Weber HM, Wang X, Midence SM. Application of the Delphi method to the development of common data elements for social drivers of health: A systematic scoping review. Transl Behav Med. 2024 Jun 27;14(7):426-433. doi: 10.1093/tbm/ibae020.

Jünger S, Payne SA, Brine J, Radbruch L, Brearley SG. Guidance on Conducting and Reporting Delphi Studies (CREDES) in palliative care: Recommendations based on a methodological systematic review. Palliat Med. 2017 Sep;31(8):684-706. doi: 10.1177/0269216317690685.

Gattrell WT, Logullo P, van Zuuren EJ, Price A, Hughes EL, Blazey P, Winchester CC, Tovey D, Goldman K, Hungin AP, Harrison N. ACCORD (ACcurate Consensus Reporting Document): A reporting guideline for consensus methods in biomedicine developed а modified Delphi. **PLoS** Med. 2024 Jan 23;21(1):e1004326. via 10.1371/journal.pmed.1004326.

Bell DR, Snedden TR, Biese KM, Nelson E, Watson AM, Brooks A, McGuine TA, Brown RL, Kliethermes SA. Consensus Definition of Sport Specialization in Youth Athletes Using a Delphi Approach. J Athl Train. 2021 Nov 1;56(11):1239-1251. doi: 10.4085/1062-6050-0725.20.

Naisola-Ruiter V. The Delphi technique: a tutorial. Research in Hospitality Management, 2022, 12(1), 91-97. doi:10.1080/22243534.2022.2080942.

Shamseer L, Moher D, Clarke M, Ghersi D, Liberati A, Petticrew M, Shekelle P, Stewart LA; PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. BMJ. 2015 Jan 2;350:g7647. doi: 10.1136/bmj.g7647.

García-Pérez MA. Are the Steps on Likert Scales Equidistant? Responses on Visual Analog Scales Allow Estimating Their Distances. Educ Psychol Meas. 2024 Feb;84(1):91-122. doi: 10.1177/00131644231164316.

Morisky DE, Ang A, Krousel-Wood M, Ward HJ. Predictive validity of a medication adherence measure in an outpatient setting. J Clin Hypertens (Greenwich). 2008 May;10(5):348-54. doi: 10.1111/j.1751-7176.2008.07572.x.

Newman-Beinart NA, Norton S, Dowling D, Gavriloff D, Vari C, Weinman JA, Godfrey EL. The development and initial psychometric evaluation of a measure assessing adherence to prescribed exercise: the Exercise Adherence Rating Scale (EARS). Physiotherapy. 2017 Jun;103(2):180-185. doi: 10.1016/j.physio.2016.11.001.

Bailey DL, Bishop A, McCray G, Foster NE, Holden MA. A new measure of exercise adherence: the ATEMPT (Adherence To Exercise for Musculoskeletal Pain Tool). Br J Sports Med. 2024 Jan 3;58(2):73-80. doi: 10.1136/bjsports-2022-106347.



Wang D, Benito PJ, Rubio-Arias JÁ, Ramos-Campo DJ, Rojo-Tirado MA. Exploring factors of adherence to weight loss interventions in population with overweight/obesity: an umbrella review. Obes Rev. 2024 May 28:e13783. doi: 10.1111/obr.13783.