



FEB 07, 2024

## Stereo-seq cell type abundance analysis

In 1 collection

Peter Kilfeather<sup>1</sup>

<sup>1</sup>University of Oxford

PK



Peter Kilfeather

OPEN ACCESS



### ABSTRACT

Stereo-seq cell type abundance analysis from Kilfeather, Khoo et al., 2024

#### DOI:

[dx.doi.org/10.17504/protocols.io.14egn3kjl5d/v1](https://dx.doi.org/10.17504/protocols.io.14egn3kjl5d/v1)

**Protocol Citation:** Peter Kilfeather 2024. Stereo-seq cell type abundance analysis. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.14egn3kjl5d/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
We use this protocol and it's working

**Created:** Feb 07, 2024

Last Modified: Feb 07, 2024

PROTOCOL integer ID: 94803

**Funders Acknowledgement:**

Aligning Science Across

Parkinson's

Grant ID: ASAP-020370

Monument Trust Discovery Award

from Parkinson's UK

Grant ID: J-1403

## Protocol

- 1 To test for differential abundance of cell types between age groups, we used mixed effects modelling of associations of single cells (MASC). MASC tests whether cell type membership of individual cells is influenced by an experimental covariate of interest, while accounting for technical covariates and biological variation. We specified mouse genotype as a fixed covariate and mouse of origin as a random effect in the generalized, mixed-effect model. Changes in cell type abundance were considered significant at  $FDR-P < 0.01$ .