



NOV 23, 2023

PCR Amplification of Clock and Adcyap1 genes with EmeraldAmp® GT PCR Master Mix in Avian species for polymorphism elucidation

Louis-Stéphane Le

Clercq^{1,2},

Desiré Lee Dalton³,

Antoinette Kotzé^{1,2}, Paul Grobler¹

¹University of the Free State;

²South African National Biodiversity Institute; ³Teesside University

OPEN ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.6qpvrw3gm/v1

External link:

<https://sites.google.com/view/lsleclercq/projects/phd-project>

Protocol Citation: Louis-Stéphane Le Clercq, Desiré Lee Dalton, Antoinette Kotzé, Paul Grobler 2023. PCR Amplification of Clock and Adcyap1 genes with EmeraldAmp® GT PCR Master Mix in Avian species for polymorphism elucidation. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.6qpvrw3gm/v1>

MANUSCRIPT CITATION:

Le Clercq, L.S., 2023.

Biological clock measures: Assessing the association between the circadian and epigenetic clock as predictors of migration phenology and biological aging in wildlife (Doctoral thesis, University of the Free State).

Biological clock measures the association between the circadian and epigenetic clock as predictors of migration and age

Tech. support email: leclercq.l.s@gmail.com



Louis-Stéphane Le Clercq

University of the Free State, South African National Biodive...

ABSTRACT

This PCR protocol is used to amplify *Clock* and *Adcyap1* gene regions in avian species which have previously shown polymorphisms, such as poly-Q runs, that correlated to migration phenology. It was tested and optimized in Woodlands kingfisher (*Halcyon senegalensis*) and Diederik cuckoos (*Chrysococcyx caprius*). The primers were designed based on those previously used by Johnson *et al.* (2007) and Steinmeyer *et al.* (2009) by comparing the relevant gene sequences for chickens (*Gallus gallus*) with several other available avian species to select primers that would account for the most common variations in primer regions, enabling more universal amplification. Individual clock gene sequences were retrieved from Genbank and aligned in BioEdit 7.2. Primers were then selected based on the annotated regions. The assay was designed using 25 µL (half) reactions of EmeraldAmp® GT PCR Master Mix, which is premixed with loading buffer for easy gel loading following PCR and does not require a long initial denaturation step (thereby shortening the run time). Gel electrophoresis was able to confirm successful amplification of a product ±280 bp long in both species. The same primers were subsequently used for sanger sequencing. A BLAST search of the resulting sequences confirmed the identity of the amplified regions.

ATTACHMENTS

[Clock_anot_Primers+Product \[BioEdit\].bio](#)

[Adcyap anot Primers+Product \[BioEdit\].bio](#)

[NZG_LeClercq_PCR_template Clock and Adcyap1 genes.xlsx](#)

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Jun 19, 2021

Last Modified: Nov 23, 2023

PROTOCOL integer ID:
50910





Keywords: Emerald Amp, Clock genes, Avian, PCR, Clock, Adcyap, Kingfisher, Cuckoo

Funders
Acknowledgement:
National Research Foundation (RSA)
Grant ID: 112062

IMAGE ATTRIBUTION


<https://www.takara-bio.com/>

GUIDELINES

- A PCR worksheet template is included for download to automatically calculate volumes.
- Check DNA template quantity and purity prior to use in PCR.
-  EmeraldAmp® GT PCR Master Mix Takara Bio Inc. Catalog #RR310A allows for direct gel loading.
- Equipment used are interchangeable with industry equivalents.
- Experiments performed at  Room temperature is always at  21 °C .
- Treated PCR products may be stored at  -20 °C until required for sequencing.
- Briefly vortex reagents and mixes as needed.






MATERIALS

Reagents:


-  EmeraldAmp® GT PCR Master Mix Takara Bio Inc. Catalog #RR310A
- **Primers:** (Inqaba Biotec. Industries; Annotated "BioEdit" alignment files are included)

A	B	C	D	E	F
Adcyap F	GATGTGAGTAACCAGCCAC T	Adcyap1	Gene ID: 408251	20	61.3
Adcyap R	ATAACACAGGAGCGGTGA	Adcyap1	Gene ID: 408251	18	59.7
Clock F1	TGGAGCAGTAATGGTACCA AGTA	clock	Gene ID: 373991	23	62.9
Clock F2	TGGAGCGGTAATGGTACCA AGTA	clock	Gene ID: 373991	23	65.0
Clock R1	TCAGCTGCGACTGAGCTGG	clock	Gene ID: 373991	19	66.0
Clock R2	TCAGCTGTGGCTGAGCTGG	clock	Gene ID: 373991	19	66.1

Summary of primer details for the assay including the primer name, sequence, gene, gene ID, length and Tm

-  UltraPure™ TBE Buffer 10X Thermo Fisher Scientific Catalog #15581044
-  SeaKem® LE Agarose Lonza Catalog #50004
-  SYBR SAFE DNA stain Life Technologies Catalog #S33102
-  Quick-Load 100 bp DNA Ladder - 375 gel lanes New England Biolabs Catalog #N0467L
-  ExoSAP-IT™ PCR Product Cleanup Reagent Thermo Fisher Scientific Australia Catalog #78201.1.ML

Equipment:

Equipment	
SimpliAmp Thermal Cycler	NAME
PCR	TYPE
Applied Biosystems	BRAND
A24811	SKU
https://www.thermofisher.com/order/catalog/product/A24811	LINK
Any standard PCR thermocycler will suffice	SPECIFICATIONS
	

Equipment	
Gel Doc XR+ Gel Documentation System	NAME
Gel Documentation System	TYPE
Bio-rad Laboratories	BRAND
1708195	SKU
https://www.bio-rad.com/en-us/product/gel-doc-xr-gel-documentation-system?ID=O494WJE8Z	LINK


Equipment	
PowerPac Basic Power Supply	NAME
Power Supply	TYPE
Bio-Rad Scientific	BRAND
1645050	SKU
https://www.bio-rad.com/en-us/sku/1645050-powerpac-basic-power-supply?ID=1645050	LINK




Samples:

- BioSample information information has been deposited to the BioProject ([PRJNA737185](https://dx.doi.org/10.17504/protocols.io.6qpvrdwk3gmk/v1)) linked to this protocol.


PROTOCOL MATERIALS

 UltraPure™ TBE Buffer, 10X Thermo
Fisher Catalog #15581044


Step 3.1

 EmeraldAmp® GT PCR Master Mix Takara Bio
Inc. Catalog #RR310A


In Guidelines, Materials, Step 1.1

 ExoSAP-IT™ PCR Product Cleanup Reagent Thermo Fisher Scientific
Australia Catalog #78201.1.ML


In Materials and [2 steps](#)

 SeaKem® LE Agarose Lonza Catalog #50004


Materials, Step 3.1

 SYBR SAFE DNA stain Invitrogen - Thermo
Fisher Catalog #S33102

In Materials and [2 steps](#)

 Quick-Load 100 bp DNA Ladder - 375 gel lanes New England
Biolabs Catalog #N0467L

Materials, Step 3.2

 UltraPure™ TBE Buffer 10X Thermo Fisher
Scientific Catalog #15581044

Materials

SAFETY WARNINGS

- ❗
 - Set up master mixes in a "DNA-free" room and laminar flow cabinet.
 - Add DNA to reaction tubes in a "DNA-loading" laminar flow cabinet.
 - Always dispose of biohazardous waste appropriately in accordance to lab regulations.
 - Always wear gloves and a lab coat.
 - Never directly look at the UV lamps.

ETHICS STATEMENT

Protocol approval for the present study was obtained from the protocol committee of the Department of Genetics, University of the Free State (approval number: Res18/2020). Ethics approvals were obtained from the University of the Free State (approval number: UFS-AED2020/0015/1709) as well as the South African National Biodiversity Institute (approval number: SANBI/RES/P2020/30). Appropriate research permits were also obtained from South African regulatory authorities including the Department of Agriculture, Land Reform, and Rural Development (Section 20 permit: 12/11/1/1/18(1824JD)).

BEFORE START INSTRUCTIONS

- Thaw reagents **On ice**.
- Wipe workspace with **10 % volume** Bleach, followed by **70 % volume** Ethanol, and ddH₂O before (and after).
- UV the relevant laminar flow cabinets.

Master Mix set-up

1 Prepare Master Mix and Samples* for PCR.


*Sample information has been deposited to BioSample and associated to the BioProject ([PRJNA737185](https://www.ncbi.nlm.nih.gov/bioproject/PRJNA737185)) which used this protocol.

(An experiment template is included in excel format.)

1.1 Set up the following Master Mix with

30m



 EmeraldAmp® GT PCR Master Mix Takara Bio Inc. Catalog #RR310A

for a **25 µL** reaction

in a DNA-free lab and laminar flow cabinet.

A	B	C	D
EmeraldAmp® GT PCR Master Mix	X2	X1	12.5
Forward primer	10 µM	0.2 µM	0.5
Reverse primer	10 µM	0.2 µM	0.5
Nuclease free water	-	-	9.5

Summary of components to add to Master Mix with the original and final concentrations as well as the relative volume in µL




EmeraldAmp GT MM



Working solutions of forward (Clk F1) and reverse (Clk R1) primers.

- 1.2 ■ Add 23 μ L Master Mix to 2 μ L DNA template* in individual thin-walled PCR tubes in a DNA-loading laminar flow cabinet. 20m



PCR reactions prepared to run thermal cycles.

*DNA isolated from avian blood samples may be highly concentrated, ensure that it is still less than 500 ng of DNA per reaction.

Thermal cycling

10s

- 2 Program and run the following thermal cycling profile on a thermal cycler, e.g. 2h



Equipment

SimpliAmp Thermal Cycler

NAME

PCR

TYPE

Applied Biosystems

BRAND

A24811









SKU

<https://www.thermofisher.com/order/catalog/product/A24811>

LINK

Any standard PCR thermocycler will suffice

SPECIFICATIONS

-  40 Cycles of:
 1. Denaturation at  98 °C for  00:00:10
 2. Annealing at  60 °C for  00:00:30
 3. Elongation at  72 °C for  00:01:00
- Infinite hold at  4 °C until ready for next steps.



Example of PCR run lasting approximately 1h30.

Electrophoresis

3 Confirm success of amplification by TAE/TBE electrophoresis.

- 3.1 Prepare a 2 % (v/v) gel with SeaKem® LE Agarose Lonza Catalog #50004 and 45mL UltraPure® TBE Buffer, 10X Thermo Fisher Catalog #15581044, pre-stained with SYBR SAFE DNA stain Life Technologies Catalog #S33102 using a casting tray and comb with sufficient wells.

Resolution capacity of different concentrations of gels.

A	B
0.3	5 to 60 kbp
0.6	1 to 20 kbp
0.7	0.8 to 10 kbp
0.9	0.5 to 7 kbp
1.2	0.4 to 6 kbp
1.5	0.2 to 3 kbp
2.0	0.1 to 2 kbp

Concentration (%) of agarose gels and their efficient range of separation in kilo base pairs.

Amount of agarose required for a small (50 mL) and large (100 mL) gel.

0.3	150mg	300mg
0.6	300mg	600mg
0.7	350mg	700mg
0.9	450mg	900mg
1.2	600mg	1.2g
1.5	750mg	1.5g
2.0	1g	2g

Concentration (%) of gels and their required amount of agarose.

*Note: SYBR SAFE DNA stain Life Technologies Catalog #S33102 is usually added at a concentration of 1 µL Stock per 10 mL TBE



SYBR Safe

3.2 Load 4 µL of PCR product to the gel alongside a molecular weight marker, e.g. Quick-Load 100 bp DNA Ladder - 375 gel lanesNew England Biolabs Catalog #N0467L, and run at 60 Volt for 00:30:00 . Possible settings for the

Equipment				
PowerPac Basic Power Supply				NAME
Power Supply				TYPE
Bio-Rad Scientific				BRAND
1645050				SKU
https://www.bio-rad.com/en-us/sku/1645050-powerpac-basic-power-supply?ID=1645050				LINK

are:

A	B
< 1kbp	5 V/cm
1-12 kbp	4-10 V/cm
>12 kbp	1-2 V/cm

Ideal voltages for resolving different size fragments.

3.3

Visualize and capture gel on an appropriate imager and paired software, e.g.

15m



Equipment

Gel Doc XR+ Gel Documentation System

NAME

Gel Documentation System

TYPE

Bio-rad Laboratories

BRAND

1708195


SKU

<https://www.bio-rad.com/en-us/product/gel-doc-xr-gel-documentation-system?ID=0494WJE8Z>

LINK

Expected result

Gel image of molecular marker and clock gene amplicons for Diederik cuckoo:



Positive amplification of clock genes viewed on a 1.5% TAE-Agarose gel.

Amplicon purification

30m

- 4 Purify the positive amplicons with

1h




ExoSAP-IT™ PCR Product Cleanup Reagent Thermo Fisher Scientific
Australia Catalog #78201.1.ML



prior to sequencing.

4.1 Mix  5 µL PCR product with  2 µL Exo-SAP IT reagent for a total volume of  7 µL . 30m




4.2 Incubate at  37 °C for  00:15:00 to degrade PCR primers and short products. 15m



4.3 Incubate at  80 °C for  00:15:00 to inactivate the 15m



 ExoSAP-IT™ PCR Product Cleanup Reagent Thermo Fisher Scientific
Australia Catalog #78201.1.ML



Exo-SAP digestion and inactivation cycles.

4.4 The PCR product is now ready for use in DNA sequencing*, SNP analyses, or other primer-extension applications.



*See Clock genes sequencing protocol (<https://protocols.io/view/abi-sanger-sequencing-of-avian-clock-genes-to-eluc-bvydn7s6>)