

Aug 29, 2024

Treatment of *D. melanogaster* with small molecules in fly food

DOI

dx.doi.org/10.17504/protocols.io.bp2l62rozgqe/v1

Natalie Kaempf¹, Patrik Verstreken¹

¹VIB-KU Leuven Center for Brain & Disease Research, 3000 Leuven, Belgium.

ASAP Collaborative Rese...



Natalie Kaempf

VIB KU Leuven

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.bp2l62rozgqe/v1

Protocol Citation: Natalie Kaempf, Patrik Verstreken 2024. Treatment of *D. melanogaster* with small molecules in fly food. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.bp2l62rozgqe/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 29, 2024

Last Modified: August 29, 2024

Protocol Integer ID: 104248

Keywords: ASAPCRN, fly food treatment, Q10, L-DOPA, R55



Funders Acknowledgement:

Aligning Science Across

Parkinson's

Grant ID: ASAP-000430

EMBO long-term postdoctoral

fellowship

Grant ID: ALTF_299-2019

Research project, FWO

Vlaanderen

Grant ID: G0A5219N

Research project, FWO

Vlaanderen

Grant ID: G0B8119N

Methusalem project

Grant ID: METH/21/05

(3M210778)

Research project, KU Leuven

Parkinson Fonds

Grant ID: EQZ-PARFON-O2010

Opening the Future grant,

Leuven Universiteitsfonds

(LUF)

Grant ID: EQZ-OPTFUP-O2010

Research project, FWO

Vlaanderen

Grant ID: G031324N

Abstract

This protocol describes the preparation of fly food with small molecules to treat flies during their aging. It can be adapted for any treatments. Details for small molecules used can be found in the "Materials" section



Materials

Coenzyme Q10, Merck, Cat# C9538-1G

- 50 mM stock solution in ethanol (99.8%), aliquoted in single-use portions and stored at -20°C
- final concentration of 1 mM with 2% ethanol

Retromer Chaperone, R55, Merck, Cat# 5310840001

- R55 (Calbiochem/Millipore) was dissolved in sterile distilled water as a 12 mM stock solution, aliquoted in single-use portions and stored at -20°C
- final concentration of 12 μ M

Levodopa (L-Dopa), Merck, Cat# PHR1271-500MG

- 15.2 mM stock in sterile distilled water on ice and stored at -80°C in single-use aliquots
- 1 mM final concentration





3,4-Dihydroxy-D-phenylalanine (D-Dopa), Merck, Cat# D9378-250MG

- 15.2 mM stock in sterile distilled water on ice and stored at -80°C in single-use aliquots
- 1 mM final concentration



Food preparation

1m

- 1 warm up the amount of fly food (standard corn meal and molasses food) required (e.g. 50ml for 45 XS vials) in the microwave, stir it in between, start with 1.5 min, mix the food, heat up for 1min again and continue until food has no clumps anymore
- 2 prepare XS vials (45 for each condition when 50ml/condition) and label them, place them in plastic boxes
- 3 measure the amount of food in measuring cylinder and let it cool down in the beaker to around  40 °C (immediately clean all glass ware after usage)
- 4 add compound/solvent and stir very well ( 00:01:00), to keep the food liquid put in water bath to maintain the temperature
- 5 pipette around  1 mL per XS vial with 10 or 25 ml pipette
- 6 Make sure the food is filling the whole bottom of the vial and shows even surface to avoid flies getting stuck in small wholes when aging
- 7 store the food in darkness at  4 °C in the cold room and clean up
- 8 use the food not longer than 2 weeks

1m

fly treatment

- 9 depending on the experimental condition flies are exposed to the treatment and corresponding control after they are collected (Q10 or R55 treatment) or 10 days prior the experiment (D-/L-DOPA treatment)
- 10 flies on treatment/control food are kept in a temperature and light-controlled incubator at 25°C and 12-hour light-dark cycle
- 11 flies are flipped to new food every two to three days



- 12 On the day of the experiment flies are flipped on fresh treatment/control food at Zeitgeber zero to one