



AUG 02, 2023

OPEN ACCESS



DOI:
dx.doi.org/10.17504/protocols.io.3byl4je2olo5/v1

Protocol Citation: Ayse Ulusoy, Michael Helwig, Angela Rollar, Michael Klinkenberg, Sinead O'Sullivan, Shirley Lee, Rita Pinto-Costa, Donato Di Monte 2023. Intravagal AAV injection in mice.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.3byl4je2olo5/v1>


License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Intravagal AAV injection in mice

Ayse Michael
Ulusoy¹, Helwig¹, Angela Rollar¹,
Michael
Klinkenberg¹, Sinead O'Sullivan¹, Shirley Lee¹,
Donato Di
Rita Pinto-Costa¹, Monte¹

¹DZNE, DI Monte Group



Ayse Ulusoy

ABSTRACT

The protocol details AAV vector injections into the mouse vagus nerve and describes pre and post-operative mouse handling.

Created: Aug 25, 2022

Last Modified: Aug 02, 2023

PROTOCOL integer ID:
69165

Keywords: ASAPCRN

- 1 Anesthetize the mouse with isoflurane and use pain killers (e.g. Temgesic) following the ethical committee procedures
- 2 Clean the surgery area with skin disinfectant, make a small incision at the midline of the neck, and isolate the left vagus nerve. Place a holder underneath to keep it isolated and to form a base to insert the needle.
- 3 Using a micropump and a 35-gauge blunt steel needle fitted to a 10-ul NanoFil syringe, inject 800 nl AAV-containing solution. Keep the needle in place for another minute to avoid backflow.
- 4 Suture and clean the wound and return the mouse to a clean cage. Follow the ethical committee's recommendations for post-operative care.

post-operative drug treatments

- 5 Post-operative drug treatments:
 - 5.1 To stimulate DREADDs, inject 1 mg/kg CNO (Tocris) dissolved in 0.9% saline intraperitoneally.

5.2 For in situ oxidative stress assessments dissolve DHE (Dihydroethidium) in saline/dimethyl sulfoxide (DMSO) (1:1 ratio) and inject intraperitoneally at a dose of 15 mg/kg.