

•



May 18, 2021

# SPARC Cat - Sham Control Chronic Cat 1, Day 14

Brett Hanzlicek<sup>1</sup>, Anna Rietsch<sup>2</sup>, Margot Damaser<sup>3</sup>

<sup>1</sup>Advanced Platform Technology Center, Louis Stokes VA Hospital, Cleveland, Ohio; <sup>2</sup>Cleveland Clinic;

<sup>3</sup>Department of Biomedical Engineering, Cleveland Clinic Lerner Research Institute, Cleveland, Ohio



#### ABSTRACT

This is a procedure for a sham control chronic cat experiment (Day 14) for cystotomy (bladder surgery). The cystotomy is performed without UroMOCA implantation. The cat is observed daily and imaged on day 0, 14 and 30 to track changes to the bladder and overall cat health in response to the cystotomy. This protocol includes urodynamics and imaging for Day 14 in the chronic experiments.

DOI

dx.doi.org/10.17504/protocols.io.bfzhjp36

PROTOCOL CITATION

Brett Hanzlicek, Anna Rietsch, Margot Damaser 2021. SPARC Cat - Sham Control Chronic Cat 1, Day 14. **protocols.io** 

https://dx.doi.org/10.17504/protocols.io.bfzhjp36

### LICENSE

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

May 05, 2020

LAST MODIFIED

May 18, 2021

PROTOCOL INTEGER ID

36617

#### MATERIALS TEXT

Cat - Domestic short-haired cat. Male or female. 6-24 months; 3.5-5.0kg

Syringe Pump - Genie Touch; Kent Scientific

Pressure Transducer - Catalog # 503067

WPI Amplifier - 4 channel transbridge

WPI National Instruments multifunction I/O device - NI USB 6259

National Instruments Laptop with Labview software

Tubing - Masterflex L/S Platinum-Cured Silicone Tubing ID 3/16 "; Cole Parmer

**BNC** cables

Tubing to catheter - APIS

Laborie Catheter - Argyl Suction Catheter, 3.5 Fr Catalog # 8890703211

Covidien Sutures - 4-0 Vicryl Polysorb (Covidien UL212)

3-0 Prolene (Ethicon 8762)

2-0 Silk Syringe

60ml Contrast

Visipaque 320mg/ml

Surgilube

Surgical instruments

BEFORE STARTING

1 week before start of experiment - Confirm absence of bladder spasm; healthy cat

12 hours before experiment, start fasting the cat

You must transport the animal chart along with the cat to surgery site. The veterinary team will record all relevant data in the chart. The chart must then go back to the housing site.

### Transport Cat

1 Transport cat from housing site to surgery site.

### Animal Prep and catheter placement

- 2 Animal is an esthetized and abdomen is shaved by the vet team. The cat is then moved into the surgery room and attached to monitors by the vet team.
- 3 Drape animal and perform betadine scrub on abdomen and genitals.
- 4 Put surgilube on 3.5Fr catheter and insert into bladder through the urethra. Advance the catheter until resistance is met, then pull back 2-3cm.
- Use gentle suction with syringe to withdraw urine from bladder through the catheter. Measure the volume and save the urine for urinalysis.
  - ■16 mL urine removed

## Cystogram - DYNA CT

6 Use 1:5 dilution of contrast to saline to visualize bladder.

Fill the bladder with 10 mL contrast and take a single CT image.

Add 5 mL more to bladder (total of 15ml) and take single CT image Take a 3D CT image with 15ml contrast:saline in the bladder **Empty Bladder** Urodynamics/Cystometry 8 Performed after cat is transferred from Isoflurane to propofol anesthetic. Fill rate is 2ml saline/minute using syringe pump. Data is recorded using Labview software. Pressure is recorded using an external pressure transducer connected to syringe pump on one end and tubing that leads to the catheter on the other end. Fill bladder with saline and record pressure using pressure transducer connected to LabView Fill to 28.6 mL; End round 1 urodynamics **Empty Bladder** 10 Urodynamics/cystometry 2 Fill bladder with saline at 2ml/min Fill to 19 mL; End round 2 urodynamics Wake and return cat Wait for cat to wake up and return to housing site. 13

