



FEB 01, 2024

🌐 Aggregation of human recombinant alpha-synuclein

Minee-Liane Choi¹

¹UCL Institute of Neurology



Minee-Liane Choi

UCL Institute of Neurology, The Francis Crick Institute

ABSTRACT

This protocol describes a method of generating aggregation of human recombinant alpha-synuclein.

OPEN  ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.6qpvr35obvmk/v1

Protocol Citation: Minee-Liane Choi 2024. Aggregation of human recombinant alpha-synuclein.

protocols.io

<https://dx.doi.org/10.17504/protocols.io.6qpvr35obvmk/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: Feb 01, 2024

Last Modified: Feb 01, 2024

PROTOCOL integer ID: 94534

Keywords: ASAPCRN

- 1 Humanrecombinant α -Syn Monomeric WT α -Syn is purified from Escherichia coli as previously described in

CITATION

Hoyer W, Antony T, Cherny D, Heim G, Jovin TM, Subramaniam V (2002). Dependence of alpha-synuclein aggregate morphology on solution conditions..

LINK

<https://doi.org/>

[Hoyer et al., 2002].

- 2 Aggregation reactions are carried out using a solution of α -Syn [M] 70 micromolar (μ M) in [M] 25 millimolar (mM) Tris buffervsupplemented with [M] 100 millimolar (mM) NaCl, pH 7.4 (in the presence of [M] 0.01 % volume % NaN3 to prevent bacterial growth).
- 3 The buffer is freshly prepared before each experiment and passed through a \pm 0.02 μ m syringe filter (Anotop, Whatman) to remove insoluble contaminants.
- 4 Prior to incubation, the reaction mixture is ultra-centrifuged at 90k r.p.m. for 01:00:00 at 4 °C to remove potential seeds. 1h
- 5 The supernatant is collected and separated in two fractions: one kept at 4 °C at all times until use(monomers), and a second incubated in the dark at 37°C 37 °C and 200 rpm , during 07:00:00 - 08:00:00 hours to avoid fibril formation (monomers+oligomers). 15h

- 6 α -Syn is always kept in LoBind microcentrifuge tubes (Eppendorf, Hamburg, Germany) to limit surface adsorption.