



Aug 29, 2024

Purification of mCherry-WIP12d/WIP13

DOI

dx.doi.org/10.17504/protocols.io.4r3l2qqyql1y/v1

Elias Adriaenssens¹, Dorotea Fracchiolla²

¹Sascha Martens lab, University of Vienna, Max Perutz Labs - Vienna;

²Sascha Martens lab, University of Vienna, Max Perutz Labs - Vienna (AT)



Elias Adriaenssens

Sascha Martens lab, University of Vienna, Max Perutz Labs - ...

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.4r3l2qqyql1y/v1**

Protocol Citation: Elias Adriaenssens, Dorotea Fracchiolla 2024. Purification of mCherry-WIP12d/WIP13. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.4r3l2qqyql1y/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 24, 2024

Last Modified: August 29, 2024

Protocol Integer ID: 101119

Keywords: ASAPCRN

**Funders Acknowledgement:****Aligning Science Across
Parkinson's (ASAP)****Grant ID: ASAP-000350****Marie Skłodowska-Curie****MSCA Postdoctoral****fellowship****Grant ID: 101062916**

Abstract

This protocol details the purification of mCherry- WIP12d/WIP13.

Materials

 Rosetta™(DE3)pLysS Competent Cells - Novagen **Merck Catalog #70956-4**

Lysis buffer:

A	B
Tris-HCl, pH 7.4	50 mM
NaCl	300 mM
MgCl ₂	2 mM
Glycerol	5%
Triton X-100	1%
Imidazole	10 mM
β-mercaptoethanol	2 mM

Wash buffer:

A	B
Tris-HCl, pH 7.4	50 mM
NaCl	300 mM
Imidazole	10 mM
β-mercaptoethanol	2 mM

SEC buffer:

A	B
Tris-HCl, pH 7.4	25 mM
NaCl	150 mM
DTT	1 mM



Purification procedure

16h 45m 30s

- 1 To purify mCherry-WIPI2d and mCherry-WIPI3, as described previously for WIPI2d (Fracchiolla et al. 2020, J Cell Biol, PMID: 32437499), fuse the coding sequence of WIPI2d or WIPI3 to a N-terminal 6xHis-TEV-mCherry-tag through cloning into a pET-DUET1 vector (available from Addgene).
- 2 After the transformation of the pET-DUET1 vector encoding 6xHis-TEV-mCherry-WIPI2d/WIPI3 in E. coli Rosetta pLysS cells (Novagen Cat# 70956-4), grow the cells in 2x Tryptone Yeast extract (TY) medium at 37 °C until an OD₆₀₀ of 0.4 and then continue at 18 °C .
- 3 Once the cells reach an OD₆₀₀ of 0.8, induce the protein expression with 100 micromolar (μM) isopropyl β-D-1-thiogalactopyranoside (IPTG) for 16:00:00 at 18 °C .
- 4 Collect the cells by centrifugation and resuspend in lysis buffer, complete EDTA-free protease inhibitors (Roche), CIP protease inhibitor (Sigma), and DNase (Sigma)).

16h

Lysis buffer:

A	B
Tris-HCl, pH 7.4	50 mM
NaCl	300 mM
MgCl ₂	2 mM
Glycerol	5%
Triton X-100	1%
Imidazole	10 mM
β-mercaptoethanol	2 mM

- 5 Sonicate cell lysates twice for 00:00:30 .
- 6 Clear the lysates by centrifugation at 18000 rpm, 4°C, 00:45:00 in a SORVAL RC6+ centrifuge with an F21S-8x50Y rotor (Thermo Scientific).

30s

45m





- 7 Filter the supernatant through an 0.45 µm filter and load onto a pre-equilibrated 5 ml His-Trap HP column (Cytiva).
- 8 After bind His tagged proteins to the column, wash the column with three column volumes of wash buffer.

Wash buffer:

A	B
Tris-HCl, pH 7.4	50 mM
NaCl	300 mM
Imidazole	10 mM
β-mercaptoethanol	2 mM

- 9 Elute the proteins with a stepwise imidazole gradient (30, 75, 100, 150, 225, 300 mM).
- 10 Pool the fractions containing the 6xHis-TEV-mCherry-WIP12d/WIP13, concentrate using a 30 kDa cut-off Amicon filter (Merck Millipore) and load onto a pre-equilibrated Superdex 200 Increase 10/300 GL column (Cytiva).
- 11 Elute the proteins with SEC buffer.

SEC buffer:

A	B
Tris-HCl, pH 7.4	25 mM
NaCl	150 mM
DTT	1 mM

- 12 Analyse the fractions by SDS-PAGE and Coomassie staining.
- 13 Pool the fractions containing purified mCherry-WIP12d or mCherry-WIP13.
- 14 After concentrating the purified protein, aliquot the protein and snap-frozen in liquid nitrogen.





15 Store the proteins at  -80 °C .