



APR 14, 2024

## 🌐 Visually-Guided Reward-Biased Behavioral Task

Raymond Murray<sup>1</sup>, Helen Schwerdt<sup>1,2</sup>

<sup>1</sup>Department of Bioengineering, University of Pittsburgh;

<sup>2</sup>Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD

ASAP Collaborative Research Network

Schwerdt Lab



Raymond Murray

Department of Bioengineering, University of Pittsburgh

### ABSTRACT

A protocol describing the visually guided, reward-biased saccade task monkeys were trained on in the original study: Schwerdt et al. 2020, *Sci. Adv.*

OPEN ACCESS



DOI:

[dx.doi.org/10.17504/protocols.io.kqdg325eev25/v1](https://dx.doi.org/10.17504/protocols.io.kqdg325eev25/v1)

**Protocol Citation:** Raymond Murray, Helen Schwerdt 2024. Visually-Guided Reward-Biased Behavioral Task. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.kqdg325eev25/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**Created:** Apr 08, 2024

**Last Modified:** Apr 14, 2024

## Behavioral Task

- 1 Monkeys performed a programmed eye movement task in which they would first fixate on a central cue displayed on a monitor in front of the them for a fixed duration chosen from a range of 1.2 - 3 s (National Institute of Mental Health, National Institutes of Health, VCortex).
- 2 The central cue would then extinguish while a second target appeared on the screen's periphery, where the monkeys would have to saccade to and fixate for another 4 s before receiving a reward.
- 3 Whether the second target appeared to the left or right of the center cue was associated with whether successful completion of a trial would result in either a big or small reward. The second target's location/reward-size association would swap after 15 to 45 trials or 50 to 100 trials.
- 4 The reward was a liquid food mixture (355 ml of Plus Nutrition Shake Vanilla, 355 ml of water, blended with two large bananas, and eight biscuits) delivered via a mouthpiece located near the animal's mouth at small and large reward volumes of 1.5 and 2.8 mL or 0.1 and 0.3 mL, respectively.
- 5 Monkeys had a large reward probability of either 50% or 25% and an intertrial interval length of a fixed 7.5 s or a fixed value of 5 to 15 s per session. Subjects would perform between 500 and 1200 individual trials per recording session.