



FEB 28, 2024

🌐 Immunoprecipitation for Flag-tagged proteins

Jean-Louis Parmasad¹

¹University of Ottawa



Jean-Louis Parmasad

DISCLAIMER

OPEN  ACCESS



DOI:

dx.doi.org/10.17504/protocols.io.q26g7poxqgwz/v1

Protocol Citation: Jean-Louis Parmasad 2024.
Immunoprecipitation for Flag-tagged proteins. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.q26g7poxqgwz/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working
We use this protocol and it's working

Created: Feb 28, 2024

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Follow institutional guidelines for ethical and safe use of biological materials.

ABSTRACT

Immunoprecipitation for Flag-tagged proteins for various biomolecular analyses.

Transfection and collection

- 1 Transfect HEK293T cells with 2µg of plasmid expressing Flag-tagged protein of interest using enhanced polyethyleneimine transfection reagent in Opti-MEM (Thermo Fisher Scientific, 31985070).
- 2 After 48h, collect cells and lyse with IP Lysis Buffer (25 mM Tris-HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA, 1% NP40, 5% Glycerol).

Prepare beads

- 3 Defrost beads on ice. Wash 3x with IP Lysis Buffer.

IP

- 4 Use 3% of the collected sample for the 'Input'.
- 5 Incubate the remaining sample with Anti-FLAG M2 Magnetic beads (Millipore Sigma, M8823) for 1 h at 4 °C in a revolving rack.
- 6 Place tubes in magnetic rack and collect 3% the supernatant as the "Flow-through" sample. Aspirate the remaining flow-through.

- 7** Wash beads with IP lysis buffer.
- 8** Add 2x Laemmli buffer (Bio-Rad, 1610747, Hercules, CA, USA) with 20% 2-mercaptoethanol (Bio-Rad, 1610710) to the beads.
- 9** Vortex, then incubate for 15 min at 75 °C to elude proteins from the beads.
- 10** Place tubes in magnetic rack and collect elution.
- 11** Run all samples on SDS-PAGE gel and visualize with method of choice.