



Aug 24, 2020

Determination of IgM concentration by the Mancini test.

Angel A Justiz-Vaillant¹¹University of the West Indies St. Augustine

1

Works for me

dx.doi.org/10.17504/protocols.io.bj9zkr76

University of the West Indies

angel.vaillant@sta.uwi.edu

Angel Justiz-Vaillant

University of the West Indies St. Augustine

DOI

dx.doi.org/10.17504/protocols.io.bj9zkr76

PROTOCOL CITATION

Angel A Justiz-Vaillant 2020. Determination of IgM concentration by the Mancini test.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bj9zkr76>

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 24, 2020

LAST MODIFIED

Aug 24, 2020

PROTOCOL INTEGER ID

40985

- 1 An appropriate anti-IgM antiserum (antibody) is poured in the center well of an agar-containing plate.
- 2 Carefully circular wells are cut and detached from the plates.
- 3 A series of standards containing known concentrations of IgM are placed in separate wells, while “unknown” human serum samples and control are placed in other wells.
- 4 A ring of precipitate forms in the area of optimal antigen-antibody concentration, meaning anti-IgM - IgM reaction as the antigen diffuses radially.
- 5 The diameters of the rings are measured and perceived normally in 48-72 hours.

- 6 Finally, a standard curve is developed using the ring diameters of the standards versus the concentrations.
- 7 A curve is then used to plot the concentration of the control and unknown IgM samples.