



JAN 29, 2024

DNA damage assessment in the adult Drosophila brain via comet assay

Mel Feany^{1,2}

¹Department of Pathology, Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts 02115;

²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815

ASAP Collaborative Research Network

OPEN ACCESS



Beatrice Weykopf

ABSTRACT

This protocol describes how to determine DNA damage in the adult drosophila brain using the comet assay

DOI:

dx.doi.org/10.17504/protocols.io.eq2lyjrrplx9/v1

Protocol Citation: Mel Feany 2024. DNA damage assessment in the adult Drosophila brain via comet assay. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.eq2lyjrrplx9/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited








Protocol status: Working
We use this protocol and it's working



Created: Jan 29, 2024






Last Modified: Jan 29, 2024


PROTOCOL integer ID: 94347


Keywords: ASAPCRN

- 1 Prepare a Lysis solution for one slide by mixing  20 mL of lysis buffer (Trevigen Comet Assay Reagent Kit) and  2 mL of DMSO
- 2 Melt LMAgarose in a boiling water bath, aliquot to 1.5ml tubes, and place at  37 °C until use
- 3 Place comet slide at  37 °C until use
- 4 Dissect 2 adult fly brains per the desired genotype in ice-cold PBS
- 4.1 Homogenize brains with a blue pestle
- 5 Combine head homogenate with  100 µL  37 °C agarose and immediately pipette  75 µL onto Comet Slide
- 6 Place slides flat at 4 °C for 20 minutes



- 7 Immerse slides in prechilled Lysis Solution  On ice for  00:45:00 45m





- 8 Immerse in a freshly prepared alkaline solution (prepare  25 mL l by mixing  0.3 g NaOH and  125 µL of 200mM EDTA in dH₂O). pH>13 for  00:30:00 at  Room temperature in the dark 30m

- 9 Drain excess buffer and wash in 1X TAE twice for  00:05:00 each 5m

- 10 Run for  00:10:00 at 23V/6mA in TAE_one volt per cm electrode to the electrode 10m

- 11 Drain excess buffer and rinse in dH₂O

- 12 Immerse slides in 70% Ethanol for  00:05:00 and then air-dry slides  Overnight 10m

- 13 The next day, prepare SYBR Green (Thermofisher) dilutant by mixing  1 µL in  10 mL of TE buffer 5m
(10 mM Tris-HCL pH 7.5, 1 mM EDTA in dH₂O) and place  50 µL on each circle of the comet assay slide
for  00:05:00 in the dark

- 14 Drain excess buffer and mount slides without DAPI in permount (Fisher Scientific)

- 15** After mounting and drying the slides, take images with 20X objective of the epifluorescence microscope and analyze the comet tails using the Image J Comet assay plugin