



APR 03, 2023

OPEN  ACCESS**DOI:**[dx.doi.org/10.17504/protocols.io.x54v9d44mg3e/v1](https://dx.doi.org/10.17504/protocols.io.x54v9d44mg3e/v1)

**Collection Citation:** michela.deleidi, Pascale Baden, María José Pérez J., Hariam Raji, Federico Bertoli 2023.

Glucocerebrosidase is imported into mitochondria and preserves complex I integrity and energy metabolism - ASAP protocol collection. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.x54v9d44mg3e/v1>

**License:** This is an open access collection distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
We use this collection and it's working

**Created:** Apr 03, 2023


**Last Modified:** Apr 03, 2023

**COLLECTION integer ID:**  
79922

## Glucocerebrosidase is imported into mitochondria and preserves complex I integrity and energy metabolism - ASAP protocol collection

michela.deleidi<sup>1</sup>, Pascale Baden<sup>1</sup>, María José Pérez J.<sup>1</sup>, Hariam Raji<sup>1</sup>, Federico Bertoli<sup>1</sup>

<sup>1</sup>German Center for Neurodegenerative Diseases (DZNE), Tübingen, 72076 Germany



Hariam Raji


### ABSTRACT

Collection of protocols of Deleidi Lab used in the publication: "Glucocerebrosidase is imported into mitochondria and preserves complex I integrity and energy metabolism"

**Keywords:** ASAPCRN,  
Glucocerebrosidase is  
imported into mitochondria  
and preserves complex I  
integrity and energy  
metabolism

FILES

Protocol



NAME

Gcase co-immunoprecipitation

VERSION 1


CREATED BY

Federico Bertoli

OPEN

→

Protocol



NAME

Single cell dissociation of brain organoids

VERSION 1


CREATED BY

Federico Bertoli

OPEN

→

Protocol



NAME

Plasmid-reprogramming of human fibroblasts

VERSION 1


CREATED BY

Federico Bertoli

OPEN

→

Protocol



NAME

Sequencing of construct

VERSION 1

CREATED BY

## Protocol



NAME

Single cell analysis of iPSC-derived midbrain organoids

**VERSION 1**

CREATED BY

Federico Bertoli

[OPEN](#) →

## Protocol



NAME

Complex I activity assay

**VERSION 1**

CREATED BY

Federico Bertoli

[OPEN](#) →

## Protocol



NAME

Midbrain-like Organoids generation from hiPSCs

**VERSION 1**

CREATED BY

Federico Bertoli

[OPEN](#) →

## Protocol



NAME

Endogenous coimmunoprecipitation

**VERSION 1**

CREATED BY

Federico Bertoli

[OPEN](#) →

## Protocol



NAME

Generation of induced pluripotent stem cells and gene correction

**VERSION 1**

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Virus production

**VERSION 1**

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Live-cell imaging

**VERSION 1**

CREATED BY

Hariam Raji

OPEN →

## Protocol



NAME

Expansion microscopy

**VERSION 1**

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Western Blot Analysis

**VERSION 1**

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Soluble and insoluble A-SYN fractionation

VERSION 1

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Mitochondrial complex activity assays

VERSION 1

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Neuromelanin staining (Fontana-Masson staining)+ TH-DAB staining on midbrain organoids

VERSION 1

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Differentiation NPCs to Dopaminergic/Midbrain Neurons

VERSION 1

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Immunofluorescence staining

VERSION 1

CREATED BY

Hariam Raji

OPEN →

## Protocol



NAME

In vitro GCase activity assay (total cell lysate)

VERSION 1

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Flag co-immunoprecipitation

VERSION 1

CREATED BY

Federico Bertoli

OPEN →

## Protocol



NAME

Constructs and generation of stable cell lines

VERSION 1

CREATED BY

Federico Bertoli

OPEN →