





Dec 03, 2021

© Geometric morphometrics nesomyinae rodent skulls V.2

Lea Terray, Christiane Denys, Steven Goodman, Voahangy Soarimalala, Aude Lalis, Raphaël Cornette¹,

christiane.denys 1

¹ISYEB MNHN





dx.doi.org/10.17504/protocols.io.bx4ppqvn

christiane.denys

Here we aim to define how skull morphology in an endemic and monophyletic clade of rodents (subfamily Nesomyinae), which show considerable morphological variation and are largely forest-dwelling, has evolved and to characterize this disparity in a phylogenetical and ecological context. We performed a two-dimensional geometric morphometric analysis on 371 dorsal and 400 ventral skull images comprising all nine extant endemic genera (26 different species) and tested the influence of ecological parameters in a phylogenetic context on size and shape.

Figure_1.pdf mada-rodent-leaterray- Supplementary_Tables_R
REVISED-protocole.docx EVISED3.LAST.docx

DOI

dx.doi.org/10.17504/protocols.io.bx4ppqvn

Lea Terray, Christiane Denys, Steven Goodman, Voahangy Soarimalala, Aude Lalis, Raphaël Cornette, christiane.denys 2021. Geometric morphometrics nesomyinae rodent skulls. **protocols.io**

https://protocols.io/view/geometric-morphometrics-nesomyinae-rodent-skulls-bx4ppqvn

christiane.denys

collection

£

Skull morphological evolution in Malagasy endemic Nesomyinae rodents

geometric morphometrics protocols, neomyine skulls, Madagascar

_____ collection,



1

Citation: Lea Terray, Christiane Denys, Steven Goodman, Voahangy Soarimalala, Aude Lalis, Raphaël Cornette, christiane.denys Geometric morphometrics nesomyinae rodent skulls https://dx.doi.org/10.17504/protocols.io.bx4ppgvn

@Christiane Denys

Sep 09, 2021

Dec 03, 2021

53103

.

Here we aim to define how skull morphology in an endemic and monophyletic clade of rodents (subfamily Nesomyinae), which show considerable morphological variation and are largely forest-dwelling, has evolved and to characterize this disparity in a phylogenetical and ecological context. We performed a two-dimensional geometric morphometric analysis on 371 dorsal and 400 ventral skull images comprising all nine extant endemic genera (26 different species) and tested the influence of ecological parameters in a phylogenetic context on size and shape.

Figure_1.pdf mada-rodent-leaterray- Supplementary_Tables_R REVISED-protocole.docx EVISED3.LAST.docx

FILES