



Aug 19, 2020

ELISA for quantification of IL-6 in human serum or plasma.

Angel A Justiz-Vaillant¹¹University of the West Indies St. Augustine
1 Works for me dx.doi.org/10.17504/protocols.io.bjy7kpzn
[University of the West Indies](#) angel.vaillant@sta.uwi.edu

Angel Justiz-Vaillant
University of the West Indies St. Augustine

ABSTRACT

Interleukin-10 is a pleiotropic cytokine playing a critical role as a regulator of myeloid and lymphoid cell function. Due to the ability of IL-10 to blocking cytokine synthesis and many accessory cell functions of antigen-presenting cells such as macrophages this cytokine is a potent suppressor of the effector functions of macrophages, T-cells and NK cells. IL-10 also participates in regulating proliferation and differentiation of B-lymphocytes, mast cells and thymocytes [1].

Reference

1. Ouyang W, O'Garra A. IL-10 Family Cytokines IL-10 and IL-22: from Basic Science to Clinical Translation. *Immunity*. 2019;50(4):871-891. doi:10.1016/j.immuni.2019.03.020

DOI

dx.doi.org/10.17504/protocols.io.bjy7kpzn

PROTOCOL CITATION

Angel A Justiz-Vaillant 2020. ELISA for quantification of IL-6 in human serum or plasma.. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bjy7kpzn>

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Aug 19, 2020

LAST MODIFIED

Aug 19, 2020

PROTOCOL INTEGER ID

40703

MATERIALS

NAME	CATALOG #	VENDOR
IL-10, Interleukin-10, human	RC212-21.SIZE.2ug	Bio Basic Inc.

- 1 An anti-human IL-6 coating antibody is adsorbed onto microwells by incubation overnight at 4°C.

- 2 Add 50 µl of human serum. Human IL-6 present in the serum sample binds to antibodies adsorbed to the microwells.
- 3 The microplate is blocked with 3% non-fat milk-PBS buffer and later washed to remove unbound proteins.
- 4 Fifty (50) µl of biotin-conjugated anti-human IL-6 antibody is added.
- 5 The microplate is rewashed with PBS-Tween buffer.
- 6 One hundred µl of streptavidin-HRP is added and binds to the biotin-conjugated anti-human IL-6 antibody.
- 7 The plate is washed following incubation to remove the unbound Streptavidin-HRP.
- 8 Add 50 µl of 3,3',5,5'- tetramethylbenzidine (TMB; Sigma-Aldrich) to each well.
- 9 Incubate the microwells in the dark for 15 min.
- 10 A colored product is formed in proportion to the quantity of human IL-6 present in the sample or standard.
- 11 The reaction is terminated by addition of 3M H₂SO₄ and absorbance is measured at 450 nm.
- 12 A standard curve is made from 7 human IL-6 standard dilutions and the human IL-6 sample concentration determined.
- 13 For better results place the microplate on a shaker in every incubation.