

Jun 19, 2024

Human RA-gastruloid induction from pluripotent stem cells

DOI

dx.doi.org/10.17504/protocols.io.261ge5epog47/v1

Nobuhiko Hamazaki¹

¹University of Washington



Nobuhiko (Nobu) Hamazaki

University of Washington

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.261ge5epog47/v1**

Protocol Citation: Nobuhiko Hamazaki 2024. Human RA-gastruloid induction from pluripotent stem cells. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.261ge5epog47/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: June 17, 2024

Last Modified: June 19, 2024

Protocol Integer ID: 101954

Keywords: Embryo model, Stem cell, Gastruloid



Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Gastruloids are a powerful in vitro model of early human development. However, although elongated and composed of all three germ layers, human gastruloids do not morphologically resemble post-implantation human embryos. Here we show an early pulse of retinoic acid (RA), together with Matrigel, robustly induces human gastruloids with posterior embryo-like morphological structures, including a neural tube flanked by segmented somites, and diverse cell types including neural crest, neural progenitors, renal progenitors, and myocytes. Through in silico staging based on single-cell RNA-seq (scRNA-seq), we find human RA-gastruloids to be more advanced than other embryo models, and comparable to E9.5 mouse and CS11 cynomolgus monkey embryos. We leverage chemical and genetic perturbations of RA-gastruloids to confirm that WNT and BMP signalling regulate somite formation and neural tube length in the human context, while transcription factors TBX6 and PAX3 underpin presomitic mesoderm and neural crest, respectively. Looking forward, RA-gastruloids are a robust, scalable model for decoding early human embryogenesis.



Materials

- StemPro Accutase (Thermo, A1110501)
- Wash media

Note

DMEM/F-12 (Thermo, 11320033) + 0.1% BSA (Thermo, 15260037)

- 10 μ M Y-27632 (Sellek, S1049)
- Nutristem hPSC XF medium (Biological Industries, 05-100-1A)
- 30mM CHIR99021 (CHIR, Millipore, SML1046)

Note

(Make 6 μ L aliquots and stored in -20C, and test the batch before the use)

- 100mM Retinoic Acid (RA) stock (Millipore Sigma, R2625)

Note

Make 10 μ L aliquots and stored in -20C)

- Essential 6 medium (Thermo, A1516401)
- Matrigel (Corning, 354230)
- Non-adhesive 96-well plate, such as Nunclon™ Sphera™ 96-Well, Nunclon Sphera-Treated,U-Shaped-Bottom Microplate and PrimeSurface® 3D culture: Ultra-low Attachment Plates: 96 well,U bottom, Clear plates
- Vitronectin (Gibco, A14700)

Note

Make 200 μ L aliquots and store them in -20C and dilute them x100 in PBS before the use and keep remaining in 4C and use it within 2 weeks



Day0, Passage human PSCs onto vitronectin-coated NutriStem

- 1 Coat wells with Vitronectin and incubate them in 37C for at least 15 min
- 2 Dissociate pre-treated ESCs
- 3 Aspirate media from wells and wash wells with PBS(-) and aspirate PBS (-)
- 4 Add 500 μ L Accutase to each well and incubate them at 37C for 4 min
- 5 Quench the reaction by adding 2 mL of Wash media containing 10 μ M Y-27632 media and pipette up and down ~10 times in a well to dissociate cells
- 6 Transfer them to a new 15mL tube and centrifuge for 3 min at 300 g
- 7 Suspend them in NutriStem containing 10 μ M Y-27632 and spread 2×10^4 cells for 1/12 well onto each vitronectin-coated well

Note

The number of seeding cells on the pre-treatment plate is important for later gastruloid elongation and needs to be optimized for each lab and each cell line.

Day 1-3, Pre-treatment of ESCs (CHIR stimulation)

- 8 **On day 1**, Change medium to fresh NutriStem medium containing 5 μ M Y-27632
- 9 **On day 2**, Change the medium to fresh NutriStem medium containing 4 μ M CHIR

**Note**

- CHIR concentrations and duration of treatment should be optimized for each CHIR batch and/or each cell line.
- CHIR should be used within three months from the aliquoting.

10 **On day 3**, Change medium to fresh NutriStem medium containing 4 μ M CHIR + 500 nM RA

Note

- The concentrations of CHIR and RA should be optimized for each cell line.

Day 4-9, Human RA-gastruloid induction

11 **On day 4**, prepare the following reagents

Note

- Accutase = 500 μ L for 1/12 well
- Wash media + 10 μ M Y-27632 = 2 mL for 1/12 well
- Essential 6 medium + 1 μ M CHIR + 5 μ M Y-27632 = 50 μ L / well + 10%

12 Aspirate media from wells and wash wells with PBS(-) and aspirate PBS (-)

13 Add 500 μ L Accutase to each well and incubate them in 37C for 4 min

14 Quench the reaction by adding 2 mL of Wash+Y media to each well and pipette up and down ~10 times in a well to dissociate cells

15 Transfer them to a new 15 mL tube and centrifuge for 3 min at 300 g

16 Wash again with 2 mL Essential6 + 1 μ M CHIR + Y and centrifuge it again



- 17 ■ Resuspend them in 1-2 mL Essential 6 + Y medium and count cells by Countess
- 18 Transfer the necessary number of cells (4000 cells/well) into a reservoir containing a medium (50 μ l/well)
- 19 Spread cells with a multi-channel pipette
- 20 Incubate them in 37C incubator
- 21 ■ **On day 5 (+24h from RA-gastruloid induction)**, add 150 μ L Essential 6 media to each well.
- 22 ■ **On day 6 (+48h from RA-gastruloid induction)**, carefully remove 150 μ L medium and add 150 μ L of fresh Essential 6 medium, containing 100 nM RA and 5% Matrigel

Note

At this moment, you would see some elongation of RA-gastruloids

- 23 **On day 9 (+120h from RA-gastruloid induction)**, you should see the fully elongated RA-gastruloid with neural tube and segmented somites