

FEB 28, 2024

# OPEN BACCESS



DOI:

dx.doi.org/10.17504/protocols.io. 36wgq3xmolk5/v1

**Protocol Citation:** sdwalto, Bryan\_Killinger, Jeffrey Kordower 2024. Rotarod. **protocols.io** https://dx.doi.org/10.17504/protoc ols.io.36wgq3xmolk5/v1

License: This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working We use this protocol and it's working

Created: Feb 28, 2024

## Rotarod

sdwalto<sup>1</sup>, Bryan\_Killinger<sup>2</sup>, Jeffrey Kordower<sup>1</sup>

<sup>1</sup>ASU NDRC; <sup>2</sup>Rush University



sdwalto

#### **ABSTRACT**

Rotarod protocol optimized for mice. This test is made to assess motor coordination and balance. The results should show lower latency to fall numbers for mice with motor impairments.

#### SAFETY WARNINGS



Depending on the rotarod apparatus, you need to be careful of mice tails being caught in-between the rotating rod and the separating wall panels.

#### **ETHICS STATEMENT**

Animal maintenance and experiments were performed in accordance with the National Institutes of Health guidelines and approved by the Institutional Animal Care and Use Committee of the Rush University Medical Center (Chicago, IL).

### protocols.io

Last Modified: Feb 28, 2024

PROTOCOL integer ID: 95890


Keywords: ASAPCRN, rotarod,

mice behavior

### **Funders Acknowledgement:**

Jeffrey Kordower

Grant ID: NIH R21 NS109871



- 2 <u>Test Period (1 hour later)</u>: Each mouse was place inside the rotarod apparatus with increasing speed (4 rpm to 40 rpm in 300 secs). Latency to fall was recorded. (Digital rotarod recorded it automatically)
- 3 Each mouse received two consecutive trials. Mean latency to fall was used for analysis.