



Jul 16, 2024

Striatal sections immunofluorescence and analysis

DOI

dx.doi.org/10.17504/protocols.io.n2bvjnpmxgk5/v1

Chuyu Chen¹

¹Northwestern University, Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815

ASAP Collaborative Rese...

Parisiadou lab



Chuyu Chen

Northwestern University, Aligning Science Across Parkinson's...

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.n2bvjnpmxgk5/v1**

Protocol Citation: Chuyu Chen 2024. Striatal sections immunofluorescence and analysis. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.n2bvjnpmxgk5/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 16, 2024

Last Modified: July 16, 2024

Protocol Integer ID: 103539

Keywords: ASAPCRN



Funders Acknowledgement:

**Aligning Science Across
Parkinson's [ASAP-020600]
through the Michael J. Fox
Foundation for Parkinson's
Research (MJFF)**

Grant ID: ASAP-020600

Abstract

Previous studies have shown that the phosphorylation of ribosomal protein S6 (p-rpS6) at S235/236 residues increases in iSPNs in response to haloperidol activation. This phosphorylation is cAMP/PKA dependent, which is the canonical pathway inhibited by D2R antagonism. We therefore explored whether LRRK2 inhibition interferes with this haloperidol-mediated phosphorylation event.

Immunostaining

- 1 Drd2-eGFP mice crossed with LRRK2-GS KI and control littermates were treated as indicated and perfused with 50 ml of PBS
- 2 Mice were perfused with 4% paraformaldehyde in PBS.
- 3 Brains were dehydrated with 30% sucrose in PBS for 48hrs and cut coronally (30 μ m) by cryostat
- 4 Striatal sections were incubated with 5% goat serum in 0.2% Triton X-100 for 2 hrs
- 5 the sections were incubated in the same solution overnight at 4 °C with the primary antibodies anti-GFP (1:1000, Invitrogen) and anti-phospho-S6 Ribosomal protein (Ser236/236) (1:300, Cell Signaling Technology)
- 6 Sections were washed with PBS for 5mins at RT
- 7 Repeat the wash with fresh PBS
- 8 Incubate with the secondary antibodies Alexa FluorTM 488 and Alexa FluorTM 647 (both 1:600, Invitrogen) for 3 hrs.
- 9 Sections were washed with PBS for 5mins at RT
- 10 Repeat the wash with fresh PBS
- 11 Sections were mounted on ProLong Diamond Antifade Mountant

Confocal imaging



12 Fluorescence images were obtained with Nikon A1R microscope and were acquired with a 20x objective at 1,024 x1,024 pixel resolution

12.1 Stitched images of the whole striatum were automatically acquired with Nikon Element

Image analysis

13 GFP and phospho-S6 Ribosomal protein signal were measured using Imaris 10.1 software

14 Surface rendering function was used to segment Drd2-eGFP cells

14.1 Background subtraction was enabled

14.2 The diameter of the largest sphere was set at 15 μm and automatically thresholded

14.3 Smooth surface was set to 1.23 μm

14.4 Segments were filtered with an area ranging between 100 μm^2 and 6,000 μm^2

15 Mean intensity of phospho-S6 Ribosomal protein within Drd2-eGFP surface above 2 times of average phospho-S6 Ribosomal protein channel mean intensity was considered positive