




Jul 08, 2022

GST Bead pulldown Assay

 Forked from [WIP12d Coprecipitation Assay](#)Adam Yokom¹¹University of California, Berkeley1 *Works for me* Share

This protocol is published without a DOI.



Adam Yokom

DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

ABSTRACT

GST Pulldown Assay for recruitment of bait proteins to GST labeled prey proteins. Prey proteins can be purified or from lysate.

PROTOCOL CITATION

Adam Yokom 2022. GST Bead pulldown Assay. **protocols.io**
<https://protocols.io/view/gst-bead-pulldown-assay-ccwksxcw>



FORK NOTE

FORK FROM

Forked from [WIP12d Coprecipitation Assay, Imstrong](#)

KEYWORDS

ASAPCRN

LICENSE

_____ This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Jul 07, 2022

LAST MODIFIED

Jul 08, 2022

PROTOCOL INTEGER ID

66220

DISCLAIMER:

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

- 1 Homogenize cell pellet with GST tagged protein in 0.5 ml of lysis buffer/protease inhibitors/1% TritonX-100. Clarify lysate by centrifugation at 40,000g for 15 min
- 2 Equilibrate 30uL of Glutathione Sepharose beads (GE Healthcare) into pulldown buffer. To do this, pipette 60uL of 50% slurry into a 1.7uL eppy. Add >500mL of wash buffer. Slow spin to pellet resin. ~1000rpm for 1 minute should be good. Repeat X3
- 3 Mix clarified lysate and washed GST resin together.
- 4 Add 1-10 μ M purified protein (add buffer of 25 mM HEPES pH 7.5, 150mM NaCl, 1mM MgCl₂, 1mM TCEP to final volume of 200 uL). Alternatively, add 10mL of HEK293GnTi lysate.

- 5 Let rock at 4C overnight. Wash x3 with buffer: 25 mM HEPES pH 7.5, 150mM NaCl, 1mM MgCl₂, 1mM TCEP
- 6 Elute washed resin with 50uL buffer: 25 mM HEPES pH 7.5, 150mM NaCl, 1mM MgCl₂, 1mM TCEP + 25 mM glutathione
- 7 Mix 17 uL eluent with 3 uL SDS-Loading dye. Heat samples for 5min @ 60C.
- 8 Run beads on SDS-PAGE gel and stain with Coomassie.