

Jun 17, 2024

Evaluation of the Antibacterial efficacy of Allium Sativum Gel as Intracanal Medicament Versus Calcium Hydroxide Paste Against Enterococcus faecalis in Single Rooted Teeth (A Comparative In-Vitro Study)

DOI

dx.doi.org/10.17504/protocols.io.36wgqndp3gk5/v1

Rania Ahmed Ahmed ELnaa¹

¹MSc student , endodontic department, faculty of dentistry cairo university



Rania Ahmed Ahmed ELnaa

MSc student , endodontic department, faculty of dentistry ca...

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.36wgqndp3gk5/v1

Protocol Citation: Rania Ahmed Ahmed ELnaa 2024. Evaluation of the Antibacterial efficacy of Allium Sativum Gel as Intracanal Medicament Versus Calcium Hydroxide Paste Against Enterococcus faecalis in Single Rooted Teeth (A Comparative In-Vitro Study).

protocols.io <https://dx.doi.org/10.17504/protocols.io.36wgqndp3gk5/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: In development

**We are still developing and
optimizing this protocol**

Created: June 13, 2024

Last Modified: June 17, 2024

Protocol Integer ID: 101766

Abstract

Objectives

The aim of this study is to evaluate the antimicrobial efficiency of garlic extract as an intracanal medicament in comparison to calcium hydroxide paste against *Enterococcus faecalis* viability developed in single rooted root canals by bacterial DNA extraction method using real time PCR.

Research Question

Will application of intracanal medicament of garlic extract differ from calcium hydroxide paste in reduction of *E. Faecalis* count when used in single rooted teeth?

Null Hypothesis

There is no difference between the antimicrobial efficacy of garlic extract and calcium hydroxide paste in reduction of *E. faecalis* count in single rooted teeth.

PICO elements

P: Population

Single rooted human freshly extracted teeth with single canals.

I: Intervention

Garlic extract used as intracanal medication.

C: Control

Calcium hydroxide paste used as intracanal medication.

O: Outcome

Bacterial load reduction determined by using real time PCR technique after root canal preparation.

Time of sampling

Sample 1 (S1): after inoculation of root canals& before instrumentation.

Sample 2 (S2): after instrumentation.

Sample 3 (S3): after intracanal medicament placement for 1 week.

Sample 4 (S4): after intracanal medicament placement for 2 weeks.

Attachments



[A2 1ry sign Rania MS...](#)

465KB

