



Sep 08, 2020

COVID19 RTLAMP Assay

matt¹, Eugene Joseph¹, Arun Manoharan Arunprimediscoveriescom¹¹Prime Discoveries

In Development

dx.doi.org/10.17504/protocols.io.bkx6kxre

Prime Discoveries

matt

DOI

dx.doi.org/10.17504/protocols.io.bkx6kxre

PROTOCOL CITATION

matt , Eugene Joseph, Arun Manoharan Arunprimediscoveriescom 2020. COVID19 RTLAMP Assay.
protocols.io
<https://dx.doi.org/10.17504/protocols.io.bkx6kxre>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Sep 07, 2020

LAST MODIFIED

Sep 08, 2020

PROTOCOL INTEGER ID

41694

MATERIALS

NAME	CATALOG #	VENDOR
RNAse and DNAse-free 1.5ml tubes		
Tubes, strips of 8	AB0452	Thermo Fisher

STEPS MATERIALS

NAME	CATALOG #	VENDOR
Binding Solution		
Wash Solution		
Resuspension Buffer		
Reaction Mix		
Primer Mix		
Positive Control		
Lysis Buffer		

EQUIPMENT

NAME	CATALOG #	VENDOR
ThermoMixer	5382000023	
Mini-centrifuge	S67601B	
SimpliAmp Thermal Cycler	A24811	

SAFETY WARNINGS

Biohazardous Materials

DISCLAIMER:

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to protocols.io is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with protocols.io, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

BEFORE STARTING

Heat incubator

Sample Lysis / Inactivation

- 1 Centrifuge the Sample in a centrifuge that can fit 15ml tubes, or let the solid debris settle for 30 min on Ice.

🕒 00:30:00

Thaw the Lysis Buffer on ice

- 2 **If using a Heat Block and Following the Full Method (Includes Purification / Concentration)**

Transfer 100ul of the sample Input Material (or clarified supernatant if Fresh Saliva) to a new 1ml tube containing an equal volume of 100ul of Lysis Buffer and pipette mix. Label this tube to indicate the sample name.



Lysis Buffer

📄 100 µl Lysis Buffer

📄 100 µl Clarified Sample

- 3 Use a tube cap lock to prevent tubes from popping open.
- 4 Place the tubes in a Heat block or water bath set at 95°C for 5 min

🕒 00:05:00



ThermoMixer
Benchtop Incubator

Eppendorf 5382000023 [↗](#)
Any heat block will suffice



- 5 Carefully transfer the tubes to ice or a 4°C Cold Block for 30 sec, or let sit at Room Temperature for 5 min.

🕒 00:00:30

🌡 4 °C

Or

🕒 00:05:00

🌡 Room temperature

- 6 Centrifuge for 3 sec at 2000xg to spin down any condensation.

🕒 00:00:03



Mini-centrifuge
Centrifuge

Fisher S67601B [↗](#)
Any standard mini centrifuge with
adapters for different tube sizes will
suffice



Samples are ready go to Step #11

- 7 **If using a 96 well Thermal Cycler and following the Quick Method (Without Purification / Concentration)**

Transfer 20ul of the sample Input Material (or clarified supernatant if Fresh Saliva) to a new 200ul tube containing an equal volume of 20ul of Lysis Buffer and pipette mix. Label this tube to indicate the sample name.




Lysis Buffer

 **20 µl Lysis Buffer**

 **20 µl Clarified Sample**

- 8 Create a Method on the Thermal Cycler for 95°C for 5 min followed by a Hold at 4°C, with the lid set to 105°C.

 **00:05:00**  **95 °C**

Hold at  **4 °C**



SimpliAmp Thermal Cycler
PCR

Applied Biosystems A24811 

Any standard PCR thermocycler will suffice



- 9 Place the tubes in the Thermal Cycler and run the method

- 10 Centrifuge for 3 sec at 2000xg to spin down any condensation.

 **00:00:03**



Mini-centrifuge
Centrifuge

Fisher S67601B 

Any standard mini centrifuge with adapters for different tube sizes will suffice



Samples are ready go to step #25

- 11 Preheat the Heat Block or Water Bath
- 12 Vortex to mix the Binding Solution so there is no sediment.



Binding Solution

- 13 To each sample tube, add 800ul of Binding Solution



Binding Solution

800 µl Binding Solution

- 14 Let the Tubes sit at Room Temperature for 10 min, every 2 min invert the tube to suspend the Binding Solution.

 00:10:00 at  Room temperature

- 15 Centrifuge for 3 sec at 2000xg to pellet the Binding Reagent in the Binding Solution Note: it should appear as a white smear at the bottom of the tube. The Binding Reagent should adhere to the bottom of the tube with a short spin on a simple benchtop centrifuge. Depending on tube type and input sample viscosity, a longer spin may be required until it is sufficiently pelleted.



Mini-centrifuge
Centrifuge

Fisher S67601B 


Any standard mini centrifuge with
adapters for different tube sizes will
suffice



- 16 Remove the supernatant with a 1ml Pipette taking care not to disturb the pellet.

 **1000 µl Supernatant**


- 17 Add 1ml of Wash Solution, wash the pellet by closing and inverting the tube several times.



 **1000 µl Wash Solution**

- 18 Centrifuge for 3 sec at 2000xg to pellet the Binding Reagent in the Binding Solution

 **00:00:03**



- 19 Remove the supernatant with a 1ml Pipette taking care not to disturb the pellet.

 **1000 µl Wash Supernatant**

- 20 Centrifuge for 3 sec at 2000xg to collect any residual Wash Solution.

 **00:00:03**



Mini-centrifuge Centrifuge

Fisher S67601B [Link](#)

Any standard mini centrifuge with adapters for different tube sizes will suffice



- 21 Remove the residual supernatant with a 200ul Pipette taking care not to disturb the pellet.

200 µl Wash Supernatant

- 22 Air dry in a clean ventilated area until the Binding Reagent is dry, it should appear opaque and not reflective, air drying should take 10-15 min, some sample types may take longer to dry, do not let drying take longer than 30 min. Note: If the lab is equipped, air drying should take place in a ventilated air cabinet to prevent cross contamination. To speed up the air drying, if available it is possible to use a Centrifuge with open tube lids, or a heated vacuum concentrator. If using these it would take less than 15 minutes as it should be evaporating less than 5-10ul of Wash Solution.

00:15:00

- 23 To each tube add 10ul of Resuspension Buffer, pipette mix to resuspend the pellet. Depending on the Sample Input, the pellet may resuspend easily or stay as clumps that will take more pipetting to break apart.



Resuspension Buffer

5 µl Resuspension Buffer

- 24 Vortex until the Binding Reagent is resuspended, you can briefly centrifuge for 3 sec at 2000xg to collect droplets from the sides of the tube. Note: As the sample contains extracted resuspended RNA, the RT-LAMP reaction should proceed the same day. The extracted samples should not be stored or frozen as it would impact the RNA integrity.

00:00:03



Mini-centrifuge Centrifuge

Fisher S67601B [Link](#)

Any standard mini centrifuge with adapters for different tube sizes will suffice



Samples are ready go to Step #25

RT-LAMP Reaction

- 25 Prepare the qPCR instrument, have Ice on hand to prepare the master Assay mix. Calculate the amount of samples being run and make sure to include the Positive Control and the H₂O.
- 26 Prepare the Assay mix by adding Reaction Mix and Primer mix according to the amount of samples being run.



Reaction Mix



Primer Mix

12 µl Reaction Mix per Sample

3 µl Primer Mix per Sample

- 27 Transfer 15ul of Assay mix into the Assay wells.

15 µl Assay Mix

- 28 Transfer 5ul of each sample into the appropriate Assay well, pipette mix.

5 µl Sample

- 29 Add the Positive Control to the Plate or Strip Tube



Positive Control

5 µl Positive Control

- 30 Transfer 5ul of H2O into the appropriate Negative Control well.

5 µl H2O

- 31 Seal the plate and run it in the qPCR instrument with the method:

Capture 2 Channels:

FAM - No Quencer = Assay Target

Cy3 - No Quencher = Control Target

 00:00:30  68.5 °C and Capture Image

Repeat 80x