



Aug 19, 2020

# Direct ELISA for investigating the binding of recombinant or chemically-made Protein-AG to immunoglobulins.

Angel A Justiz-Vaillant<sup>1</sup><sup>1</sup>University of the West Indies St. Augustine**1** Works for me [dx.doi.org/10.17504/protocols.io.bjxwkppe](https://dx.doi.org/10.17504/protocols.io.bjxwkppe)

University of the West Indies

[angel.vaillant@sta.uwi.edu](mailto:angel.vaillant@sta.uwi.edu)Angel Justiz-Vaillant  
University of the West Indies St. Augustine

## ABSTRACT

Protein AG (SpAG) is an immunoglobulin-binding protein that interacts with the Fc regions of many mammalian immunoglobulins [1].

## References

1. Vaillant AJ, McFarlane-Anderson N, Wisdom B, Mohammed W, Vuma S, et al. (2013) Immunoglobulin-binding Bacterial Proteins (IBP) Conjugates and their Reactivity with Immunoglobulin in Enzyme-Linked Immunosorbent Assays (ELISA). J Anal Bioanal Tech 4: 175. doi:10.4172/2155-9872.1000175

## DOI

[dx.doi.org/10.17504/protocols.io.bjxwkppe](https://dx.doi.org/10.17504/protocols.io.bjxwkppe)

## PROTOCOL CITATION

Angel A Justiz-Vaillant 2020. Direct ELISA for investigating the binding of recombinant or chemically-made Protein-AG to immunoglobulins.. **protocols.io**  
<https://dx.doi.org/10.17504/protocols.io.bjxwkppe>

## LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

## CREATED

Aug 19, 2020

## LAST MODIFIED

Aug 19, 2020

## PROTOCOL INTEGER ID

40662

## MATERIALS

NAME	CATALOG #	VENDOR
Nunc®; 96-Well Polystyrene Round Bottom Microwell Plates, V 96 well plate, Non-Treated, clear, without lid, Sterile	260210	Thermo Fisher
Protein-L from P. Magnus		
Streptococcal protein G by Sigma Aldrich		

**1** This ELISA is used to study the interaction of protein-AG (SpAG) with diverse immunoglobulins.

- 2 The 96 well microtitre plate is coated overnight at 4°C with 1 µg/µl per well of purified immunoglobulins or 50 µl of any animal sera in carbonate-bicarbonate buffer pH 9.6.
- 3 Then plate is treated with bovine serum albumin solution and washed 4X with PBS-Tween.
- 4 Then 50 µl of peroxidase-labeled-protein-AG conjugate diluted 1:3000 in PBS-non-fat milk is added to each well and incubated for 1.30h at RT. After that the plate is washed 4X with PBS-Tween.
- 5 Pipette 50 µl of 3,3',5,5' - tetramethylbenzidine (TMB; Sigma-Aldrich) to each well.
- 6 The reaction is stopped with 50 µl of 3M H<sub>2</sub>SO<sub>4</sub> solution.
- 7 The plate is visually assessed for the development of colour and read in a microplate reader at 450 nm.
- 8 A cut-off point should be calculated as the mean of the optical density of negative controls x 2 SD.