



## May 04, 2022

## Linker Design and Optimization

## boc.protac 1

<sup>1</sup>BOC Sciences

1



dx.doi.org/10.17504/protocols.io.yxmvmnw5bg3p/v1

**PROTAC** 

boc.protac

The linker is responsible for linking two key domains of PROTAC: the domain that specifically binds to the target protein that needs to be degraded, and the domain that binds to the E3 ubiquitin ligase. To date, several types of connectors have been reported and used in the formation of PROTAC ternary complexes, such as PEG liner, alkyl liner and "click chemical" liner. It plays an important role in the effective ubiquitination and final degradation of target proteins.

As a leading service provider in drug discovery and research, BOC Sciences is fully qualified and committed to providing one-stop PROTAC® development, which has become a promising strategy in the field of small molecular drug discovery. With a comprehensive and advanced platform, we provide <u>Linker Design and Optimization</u> Services to customers around the world to meet new drug discovery goals.

DOI

dx.doi.org/10.17504/protocols.io.yxmvmnw5bg3p/v1

https://ptc.bocsci.com/services/linker-design-and-optimization-services.html

boc.protac 2022. Linker Design and Optimization. **protocols.io** https://dx.doi.org/10.17504/protocols.io.yxmvmnw5bg3p/v1

protocol ,

May 04, 2022

May 04, 2022

61936



1

Citation: boc.protac Linker Design and Optimization <a href="https://dx.doi.org/10.17504/protocols.io.yxmvmnw5bg3p/v1">https://dx.doi.org/10.17504/protocols.io.yxmvmnw5bg3p/v1</a>