



Version 3

May 11, 2021

SYBR green RT-PCR assay for the surveillance of SARS-CoV-2 variants of concern V.3

Pedro Cardoso¹, Fernando do Couto Motta², Marilda Agudo Mendonça Teixeira de Siqueira², Daniela Tupy de Godoy¹, Rodrigo Brindeiro³, Monica Barcellos Arruda¹, Elisabete Andrade¹, Marisa Ribeiro¹, Marcela Fontana-Maurell¹, Elaine Costa¹, Daniele Rocha¹, Patricia Alvarez¹

¹Immunobiological Technology Institute (Biomanguinhos);

²Laboratório de Vírus Respiratórios e Sarampo do Instituto Oswaldo Cruz (IOC/Fiocruz); ³LAVIMOAN - UFRJ

2

Works for me

dx.doi.org/10.17504/protocols.io.butznwp6

Coronavirus Method Development Community LATED

Patricia Alvarez

Immunobiological Technology Institute (Biomanguinhos)

ABSTRACT

The emergence of SARS-CoV-2 variants with multiple shared mutations has been described to be more transmissible and could affect COVID-19 morbidity and mortality. Some of these variants, known as B.1.1.7 (originally described in the United Kingdom), B.1.351 (originally described in South Africa), and P.1 (originally described in Brazil) have rapidly become dominant within their countries and require a vigorous public health response. Whole Genomic sequencing remains the gold standard method to identify the SARS-CoV-2 variant, even though this approach is laborious, time-consuming, and expensive. Here, we have developed a fast and simple SYBR green-based real-time RT-PCR assay that identifies a distinct signature affecting the Non-Structural Protein 6 (NSP6), a nine-nucleotides deletion leading to amino acid losses: $\Delta 106S$, $\Delta 107G$, and $\Delta 108F$. This NSP6 signature is present in the variants of concern (VOC) described above (B.1.1.7; B.1.351; P.1) and has not been detected in other SARS-CoV-2 lineages. The new version has corrections made on the master mix and authorship.

DOI

dx.doi.org/10.17504/protocols.io.butznwp6

PROTOCOL CITATION

Pedro Cardoso, Fernando do Couto Motta, Marilda Agudo Mendonça Teixeira de Siqueira, Daniela Tupy de Godoy, Rodrigo Brindeiro, Monica Barcellos Arruda, Elisabete Andrade, Marisa Ribeiro, Marcela Fontana-Maurell, Elaine Costa, Daniele Rocha, Patricia Alvarez 2021. SYBR green RT-PCR assay for the surveillance of SARS-CoV-2 variants of concern. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.butznwp6>
Version created by Daniela Tupy de Godoy

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

May 07, 2021

LAST MODIFIED

May 11, 2021

PROTOCOL INTEGER ID

49753

MATERIALS TEXT

- Go Taq 1-step RT-qPCR System (Promega) A2060

 [GoTaq OneStep RT qPCR System Quick Protocol FB127.pdf](#)

- Nuclease-free water
- Primers

A	B
	NSP6: ΔS106; ΔG107 e ΔF108
Primer name	Sequence
Forward	5'-TGGTTGGATATGGTTGATACTAGTT-3'
Reverse	5'-AGCTGATGCATACATAACACAGT-3'

Primers details

- Positive controls: (1) SARS-CoV-2 RNA from virus culture of the Brazilian variant P2 and (2) SARS-CoV-2 RNA from virus culture of the Brazilian variant P1. The SARS-CoV-2 RNA was kindly provided by the Respiratory Virus and Measles Laboratory – IOC (LVRS-IOC).
- Applied Biosystems™ 7500 Real-Time PCR Systems (Thermo Fischer Scientific)
-


DISCLAIMER:

This SYBR green RT-PCR protocol is for research purposes only. It should not be used for clinical diagnosis. The intention of this assay is to screen for the presence of a distinct signature affecting the Non-Structural Protein 6 (NSP6) ΔS106; ΔG107 e ΔF108, present in high-virulent SARS-CoV-2 variants

- 1 Briefly vortex and centrifuge reagents before use.
- 2 Prepare 10 μM working stocks of the primers.
- 3 Thaw the GoTaq® qPCR Master Mix (Promega), Nuclease-Free Water, and primers working stocks.
- 4 Use the 10 μM working stocks to prepare the mix, containing:

A	B	C	D
	Volume (per reaction)	final concentration	
GoTaq® qPCR Master Mix, 2X	7.5 µl	1x	
GoScript™ RT Mix for 1-Step RT-qPCR (50X)	0.3 µl	1x	
Primer Forward	0.15 µl	0.1 µM	
Primer Reverse	0.15 µl	0.1 µM	
Nuclease-free water	1.9 µl		

- 5 Add 10µl to each well or PCR tube.
- 6 Add 5 µl of RNA from positive controls 1 and positive control 2 to the correspondent well or PCR tube. Add 5 µl nuclease-free water to the negative control well or PCR tube. Mix by pipetting.
- 7 Add 5µl of extracted RNA (unknown sample) to the designated wells and mix by pipetting.
- 8 Seal plates and tubes.
- 9 Run the PCR with the following cycler conditions



7500 Real Time PCR System
ABI 4351104

