



Apr 18, 2022

Rotarod assay to detect motor phenotypes in mice

William Hancock-Cerutti^{1,2,3,4,5}, Pietro De Camilli^{1,2,3,5}

¹Departments of Neuroscience and of Cell Biology, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

²Howard Hughes Medical Institute;

³Program in Cellular Neuroscience, Neurodegeneration and Repair, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

⁴Interdisciplinary Neuroscience Program and MD-PhD Program, Yale University School of Medicine, New Haven, Connecticut 06510, USA;

⁵Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815

1



dx.doi.org/10.17504/protocols.io.j8nlkdd1dl5r/v1

 William Hancock-Cerutti

This protocol describes an assay to detect motor deficits in mice by testing how long they can stay on a rotating rod before falling.

[HC-Protocol-Rotarod.docx](#)

DOI

dx.doi.org/10.17504/protocols.io.j8nlkdd1dl5r/v1

William Hancock-Cerutti, Pietro De Camilli 2022. Rotarod assay to detect motor phenotypes in mice. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.j8nlkdd1dl5r/v1>



protocol ,

Apr 18, 2022

Apr 18, 2022

60897