


Jun 18, 2024 Version 1

🌐 Fixation and immunocytochemistry (fluorescence) in PFF-treated cultures V.1

DOI

dx.doi.org/10.17504/protocols.io.x54v922q4l3e/v1



Maria Iuliano^{1,2}

¹Yale University; ²Tufts University

ASAP Collaborative Rese...

Maria's Workspace



Maria Iuliano

Biederer Lab, Yale University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.x54v922q4l3e/v1

Protocol Citation: Maria Iuliano 2024. Fixation and immunocytochemistry (fluorescence) in PFF-treated cultures. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.x54v922q4l3e/v1> Version created by **Maria Iuliano**

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: June 06, 2024

Last Modified: June 18, 2024

Protocol Integer ID: 101845

Keywords: ASAPCRN

**Funders Acknowledgement:****ASAP**

Grant ID: 020616

Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

This protocol outlines methods used for the preparation of in vitro models for fluorescent imaging including primary rat and mouse hippocampal cultures treated with PFFs.

Image Attribution

Maria Iuliano -60x image of rat hippocampal neurons (DIV28) after 7-day treatment with PFFs. Map2 (Blue) and phospho-a-syn Ser129 (green)

Guidelines

Parameters used in this protocol may be optimized and modified for individual use depending on experimental needs.

Materials

1% SDS for PFF inactivation
4% PFA 4% sucrose in 1x PBS
1x PBS
ICC Buffer: 5% NGS in 1x PBS
ICC Blocking Buffer: 5% NGS 0.3% Triton-X in 1x PBS\
Primary & secondary antibodies of choice
Microscope slides
Forceps
Mounting medium of choice





Protocol materials

 Aqua-Poly/Mount **Polysciences, Inc. Catalog #18606-100** Step 9

Safety warnings

- ⚠ When working with PFFs, it is best to follow proper safety precautions and wear appropriate PPE, including lab coat, gloves, sleeve guards, safety glasses or face shield, and N95 or FFP2 mask.

Inactivation and disposal of PFFs and PFF-contaminated items can be done using 1% SDS as described in literature.

Collect and dilute solutions into 10vol 1% SDS for  01:00:00 at  Room temperature

Discard used tips into primary collection container with 1% SDS

Inactivated liquid can be discarded with liquid waste

Discard inactivated items in primary container into biohazard

Wipe surfaces or contaminated tools down completely with 1% SDS, followed by water, then EtOH

CITATION

Bousset L, Brundin P, Böckmann A, Meier B, Melki R (2016). An Efficient Procedure for Removal and Inactivation of Alpha-Synuclein Assemblies from Laboratory Materials..

LINK

<https://doi.org/10.3233/JPD-150691>

Before start

Please read safety guidelines ahead of working with PFFs



Fixation and Immunocytochemistry (Fluorescence)


4h 30m

- 1 In a cell culture hood remove media and wash briefly with 1x PBS.

2m

Safety information

Collect media and first wash as PFF waste for inactivation. Once in fixative, plate(s) can be moved to benchtop.

- 2  20 rpm, Room temperature , 00:15:00 Fix using 4% PFA 4% sucrose in 1xPBS with gentle rocking/shaking.

15m

Note

Check whether the antibodies you intend to use require specific fixation. Optimization may be needed.


- 3  20-30 rpm, Room temperature , 00:10:00 Wash 1x PBS. Repeat 3 times.

30m



Note

Pause Point: Coverslips can be stored after fixation for up to 2 weeks (or maybe more), though immediate processing is ideal. Store at 4C away from light, with the plate wrapped in parafilm to reduce evaporation.

II

- 4  20-30 rpm, Room temperature , 00:45:00 Block in blocking buffer (5% NGS, 0.3% Triton-X in 1x PBS)

45m

- 5  20 rpm, 4°C  Overnight Incubate in solution containing primary antibodies diluted in 5% NGS in 1x PBS



**Note****Antibodies**

A	B	C
Antibody	Company	Catalog #
Gephyrin	Synaptic Systems	147111
HA (HA-tag)	BioLegend	901513
HA (HA-tag)	Cell Signaling	3724
Homer	Synaptic Systems	160003
MAP2	Millipore	AB5543
MAP2	Millipore	MAB364
Neurofilament	Millipore	AB5539
Phospho- α -syn (Ser129)	Biolegend	825701
VGAT	Synaptic Systems	131003
VGLUT1	Millipore	AB5905

Some commonly used antibodies in our lab for PFF-treated cultures. Dilutions may need to be optimized to individual needs.

6 20-30 rpm, Room temperature , 00:10:00 Wash 1x PBS. Repeat 3 times.

30m

7 20 rpm, 4°C, 02:00:00 Incubate in solution containing secondary antibodies diluted in 5% NGS in 1x PBS

2h

Note

Secondary incubation can vary (up to O/N), but typically 2-3hr at 4C works well. 1hr at RT can also work, though it may result in excess background signal, depending on the antibody

Typically, we use Alexa Fluor secondaries from Life Technologies, except for anti-chicken CF 405 (Millipore SAB4600466).

8 20-30 rpm, Room temperature , 00:10:00 Wash 1x PBS. Repeat 3 times.

30m



9 Mount coverslips onto microscope slides using




Aqua-Poly/Mount **Polysciences, Inc. Catalog #18606-100**

or preferred mounting

medium.

5m

10 Let cure  48:00:00 before imaging

2d

Citations

Bousset L, Brundin P, Böckmann A, Meier B, Melki R. An Efficient Procedure for Removal and Inactivation of Alpha-Synuclein Assemblies from Laboratory Materials.

<https://doi.org/10.3233/JPD-150691>