




Sep 15, 2022

Coating coverslips for cell culture

Chuyu Chen¹, Ciarra Smith¹¹Northwestern university

1 Works for me

 Sharedx.doi.org/10.17504/protocols.io.5qpvrnd9v4o/v1 divya.darwinarulseeli

ABSTRACT

Coated cover-slips provide a nourishing adherent surface for cell culture. This protocol provides step by step instruction on how to coat coverslips for cell culture.

DOI

dx.doi.org/10.17504/protocols.io.5qpvrnd9v4o/v1

PROTOCOL CITATION

Chuyu Chen, Ciarra Smith 2022. Coating coverslips for cell culture. **protocols.io**
<https://protocols.io/view/coating-coverslips-for-cell-culture-cgpgtvjw>



LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Sep 15, 2022

LAST MODIFIED

Sep 15, 2022

PROTOCOL INTEGER ID

70088

Poly L-Lysine/ Poly D-Lysine coating

- 1 Precoat the coverslips with poly-L-lysine (P5899) or poly-D-lysine (P7280) at 100 µg/ml.
- 2 This is incubated for 30 minutes or more at room temperature. The coverslips are washed

with sterile water and allowed to dry.

- 3 They can be stored at room temperature until ready to use (up to 1 month).

Day before culturing: Laminin coating

- 4 Coverslips are then coated with 30 - 50 μ l of laminin (50 μ g/ml) which has been warmed. This is prepared the day before culturing of cells.
- 5 Incubate overnight in a humidified 37°C, 5% CO₂ incubator or at 37 °C for 30 minutes.
- 6 Remove excess laminin and rinse with PBS twice. Alternatively, wash with DME (50 μ L) twice.
- 7 Add the cells to the coverslips/slides and culture as needed.