




Oct 08, 2022

Behavioral tests in rodents coupled with dopamine signaling manipulations

Chuyu Chen¹, Loukia Parisiadou¹¹Northwestern university

1 Works for me

 Sharedx.doi.org/10.17504/protocols.io.261ge345jl47/v1 divya.darwinarulseeli

ABSTRACT

These two protocols describe behavioral tests associated with Parkinson's disease phenotypes in mice. The striatal motor learning protocol (rotarod test) is a two-phase paradigm assessing striatal motor learning under reversible dopamine antagonism.

DOI

dx.doi.org/10.17504/protocols.io.261ge345jl47/v1

PROTOCOL CITATION

Chuyu Chen, Loukia Parisiadou 2022. Behavioral tests in rodents coupled with dopamine signaling manipulations. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.261ge345jl47/v1>



FUNDERS ACKNOWLEDGEMENT

Aligning Science Across Parkinson's through the Michael J. Fox Foundation for Parkinson's Research (MJFF)
Grant ID: ASAP-020600

KEYWORDS

Motor function, behavior, rota-rod, open-field

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 08, 2022

LAST MODIFIED

Oct 08, 2022

PROTOCOL INTEGER ID

71030

MATERIALS TEXT




Figure 1: Schematic of the rotarod training paradigm including reversal dopamine receptors antagonism.

Rotarod test

- 1 General Rota-rod test is performed as follows:
 - 1.1 The rotarod apparatus (Panlab LE8205, Harvard Apparatus) equipped with a mouse rod (3 cm diameter)
 - 1.2 Set the rotarod with a start speed of 4 rpm, and set to 4–40 rpm acceleration over 300s
 - 1.3 Hold the mouse by tail. Place the mouse on the rod and face away from the direction of rotation.
 - 1.4 Start acceleration and record the time for each mouse to fall.
 - 1.5 Perform five trials with intertrial-interval = 15 s, max trial duration = 300 s (one daily session)

1.6 Calculate the average time to fall over five trials.

Details of dopamine signaling manipulations for Rota-rod

- 2 The specific task consists of eighteen daily sessions divided into two phases (Figure 1).
- 3 Administer intraperitoneally dopamine receptor antagonists for sessions 1-5 30 min prior to testing prepared in 0.9% saline): cocktail of 1 mg/kg SCH23390 + 1 mg/kg eticlopride, 1 mg/kg SCH23390, or 1 mg/kg eticlopride.
- 4 Following a 72-hr break, test the mice for another thirteen sessions (drug-free recovery phase)

Open field test

- 5 The open field arena (56 x 56 cm) is used to assess locomotor activity.
- 6 Turn on the lights in the recording chamber. Clean the arena with 70% ethanol and adjust the camera border to match the arena edge.
- 7 Wait for 5min to allow the ethanol to evaporate completely prior test.
- 8 A mouse is placed in the center of the arena and its ambulation activity is collected by the LimeLight software for 20 minutes.
- 9 Clean the arena with 70% ethanol between tests.
- 10 The LimeLight software provides the total distance traveled and the percentage of time/distance within different parts of the arena.

