

Sep 22, 2021

Determine the Size of Sonicated Samples and the DNA Concentration

In 1 collection

Vasso Makrantonis¹, Daniel Robertson¹, Adele L. Marston¹¹The Wellcome Centre for Cell Biology, Institute of Cell Biology, School of Biological Sciences, University of Edinburgh, Edinburgh, UK

1 Works for me

Share

dx.doi.org/10.17504/protocols.io.bn33mgqn

Springer Nature Books

Satyavati Kharde
Springer Nature

ABSTRACT

A plethora of biological processes like gene transcription, DNA replication, DNA recombination, and chromosome segregation are mediated through protein–DNA interactions. A powerful method for investigating proteins within a native chromatin environment in the cell is chromatin immunoprecipitation (ChIP). Combined with the recent technological advancement in next generation sequencing, the ChIP assay can map the exact binding sites of a protein of interest across the entire genome. Here we describe a step-by-step protocol for ChIP followed by library preparation for ChIP-seq from yeast cells.

DOI

dx.doi.org/10.17504/protocols.io.bn33mgqn

EXTERNAL LINK

https://link.springer.com/protocol/10.1007/978-1-4939-9520-2_10

PROTOCOL CITATION

Vasso Makrantonis, Daniel Robertson, Adele L. Marston 2021. Determine the Size of Sonicated Samples and the DNA Concentration. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bn33mgqn>

MANUSCRIPT CITATION please remember to cite the following publication along with this protocol

Makrantonis V., Robertson D., Marston A.L. (2019) Analysis of the Chromosomal Localization of Yeast SMC Complexes by Chromatin Immunoprecipitation. In: Badrinarayan A. (eds) SMC Complexes. Methods in Molecular Biology, vol 2004. Humana, New York, NY. https://doi.org/10.1007/978-1-4939-9520-2_10

COLLECTIONS ⓘ

**Analysis of the Chromosomal Localization of Yeast SMC Complexes by Chromatin Immunoprecipitation**

KEYWORDS

Chromatin immunoprecipitation, *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, Cohesin, Condensin, Mitosis, Meiosis, Scc1, Rec8, Brn1

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited






CREATED

Oct 27, 2020

LAST MODIFIED

Sep 22, 2021

OWNERSHIP HISTORY

Oct 27, 2020	 Lenny Teytelman	protocols.io
Jul 05, 2021	 Emma Ganley	protocols.io
Aug 24, 2021	 Satya K	
Aug 26, 2021	 Satyavati Kharde	Springer Nature
Aug 26, 2021	 Satyavati Kharde	Springer Nature

PROTOCOL INTEGER ID

43867

PARENT PROTOCOLS

Part of collection

[Analysis of the Chromosomal Localization of Yeast SMC Complexes by Chromatin Immunoprecipitation](#)

GUIDELINES

Chromatin immunoprecipitation (ChIP) is broadly used to study chromatin dynamics. Changes in occupancy of chromosomal proteins at specific loci within the genome can be measured by using ChIP-qPCR. However, this technique is costly and time consuming with high variability per experiment. Alternatively, ChIP-seq can be used to measure differences in a protein's occupancy genome wide. Finally, calibrated ChIP-seq is essential when measuring changes in occupancy between different experimental samples.

Here we describe an optimized ChIP protocol for yeast SMC proteins that can be completed within 3 days for samples analyzed by qPCR and 4 days for samples to be further processed by calibrated deep sequencing. The protocol encompasses five distinct steps: cross-linking and cell harvesting; cell lysis and sonication; immunoprecipitation, decross-linking and DNA extraction and finally determination of the size and DNA concentration of sonicated samples. These five steps are outlined here.

Yeast Strains and Growth Material:

1. Haploid *S. cerevisiae* strains of w303 background we have used include: (a) no tag control (AM1176), (b) *SCC1-6HA* (AM1145), (c) *BRN1-6HA* (AM5708), (d) *SCC2-6HIS-3FLAG* (AM6006), and (e) *SCC1-6HA pMET3-CDC20* (AM1105) as previously described [9,10,11,12].
2. For studies of protein occupancy during meiosis we have used diploid *S. cerevisiae* strains of SK1 background including (a) *REC8-3HA ndt80Δ* (AM4015), as previously described [13] and (b) *REC8-6HIS-3FLAG* (AM11000).
3. Haploid *S. pombe* strains used for calibration are: (a) *RAD21-3HA* (spAM76), (b) *RAD21-6HA* (spAM635), (c) *RAD21-6HIS-3FLAG* (spAM1863), or (d) *CND2-6HA* (spAM1862).
4. YPDA media: 1% yeast extract, 2% peptone, 2% glucose.
5. YPG agar plates: 1% yeast extract, 2% peptone, 2.5% glycerol, 2% agar.
6. YPDA4% agar plates: 1% yeast extract, 2% peptone, 4% glucose, 2% agar.
7. BYTA media: 1% yeast extract, 2% Bacto tryptone, 1% potassium acetate, 50 mM potassium phthalate.
8. SPO media: 0.3% potassium acetate, pH 7.0.
9. YES media: 0.5% yeast extract, 3% glucose, 225 mg/L supplements.

Equipment and Reagents:

1. 37% formaldehyde solution for molecular biology.
2. 2.5 M glycine: Dissolve 93.8 g glycine in ddH₂O (may require gentle heating) and bring up to 500 ml with ddH₂O.
3. Diluent buffer: 0.143 M NaCl, 1.43 mM EDTA, 71.43 mM Hepes–KOH pH 7.5.
4. TBS buffer: 20 mM Tris–HCl pH 7.5, 150 mM NaCl.
5. 2× FA lysis buffer: 100 mM Hepes–KOH pH 7.5, 300 mM NaCl, 2 mM EDTA, 2% Triton X-100, 0.2% Na-deoxycholate.
6. FastPrep screw-cap tubes.
7. 100 mM PMSF.
8. Protease inhibitor tablets Complete EDTA free.
9. Zirconia/Silica beads 0.5 mm diameter.
10. FastPrep-24 5G Homogenizer.
11. Bioruptor Twin.
12. Dynabeads Protein G.
13. Magnetic rack.
14. ChIP Wash buffer 1—low salt: 1× FA lysis buffer, 0.1% SDS, 275 mM NaCl.
15. ChIP Wash buffer 2—high salt: 1× FA lysis buffer, 0.1% SDS, 500 mM NaCl.
16. ChIP Wash buffer 3: 10 mM Tris–HCl pH 8.0, 0.25 M LiCl, 1 mM EDTA, 0.5% NP-40, 0.5% Na-deoxycholate.
17. ChIP Wash buffer 4 (TE): 10 mM Tris–HCl pH 8.0, 1 mM EDTA.
18. Chelex 100 Resin.
19. 10 mg/ml Proteinase K
20. TES buffer: 50 mM Tris–HCl pH 7.5, 10 mM EDTA, 1% SDS.
21. Nuclease-free molecular biology grade water.
22. Filter tips.
23. Luna Universal Probe qPCR Master Mix.
24. LightCycler 480 Multiwell Plate 96.
25. LightCycler real-time PCR.
26. Qiagen purification kit.
27. LoBind DNA microcentrifuge tubes.
28. Quick blunting kit.
29. AMPure XP beads.
30. Klenow 3' to 5' exo minus.
31. Quick ligation kit (T4 DNA ligase).
32. NEXTflex DNA Barcodes—12 (Bioo Scientific; #NOVA-514102).
33. Phusion High-Fidelity DNA polymerase.
34. DynaMag-PCR magnet.
35. WizardSV Gel and PCR cleanup system.
36. Qubit dsDNA-HS Assay kit (Invitrogen).
37. Qubit Fluorometric Quantitation machine.
38. Agilent 2100 Bioanalyzer system.
39. High Sensitivity DNA Reagents kit (Agilent Technologies).
40. High Sensitivity DNA Chips (Agilent Technologies).
41. MiniSeq High throughput Reagent Kit (150-cycle) (Illumina).
42. Illumina Mini-seq.

SAFETY WARNINGS

For hazard information and safety warnings, please refer to the SDS (Safety Data Sheet).

Formaldehyde and PMSF are toxic if inhaled, ingested or absorbed through the skin. Always wear a lab coat and gloves, and work in a chemical hood.

BEFORE STARTING

Use sonicated chromatin samples (step 15 from "Cell Lysis and Sonication") to determine the fragment size.

Determine the Size of Sonicated Samples and the DNA Concentration

3h 0m 30s

1 

To a **100 µl Input sample** add **80 µl TE buffer** containing **300 Millimolar (mM) NaCl**.

2 

30s

Decross-link at **65 °C** **Overnight**.

3 

1h

Add **2 µl RNase A (10 mg/ml)** and incubate at **37 °C** for **01:00:00**.

RNase treatment is important as high levels of RNA will interfere with DNA purification when using commercially available PCR purification kits. DNA yield can be markedly reduced as the columns become saturated.

4 

2h

Add **20 µl Proteinase K (10 mg/ml)** and incubate at **65 °C** for **02:00:00**.

5 

Purify DNA using a PCR purification kit. Run purified DNA on a 2% agarose gel with a 100 bp DNA ladder marker to determine fragment size. Ideally sonication should yield an enrichment of fragments between 200 and 400 bp (Fig. 2a).




Fig. 2

(a) Representative image of mitotic yeast cells sonicated with a Bioruptor Twin (Diagenode) for a 30-min round (power setting: High, 30 s ON/30 s OFF). DNA from two different samples was loaded on a 2% agarose gel with a 100 bp marker ladder.
(b) Representative optimal BioAnalyzer trace (upper panel) and contaminated trace with adapter (bottom panel)
(c). Examples of profiles generated by chromatin immunoprecipitation followed by sequencing (ChIP-seq) of the cohesin subunit Scc1 in wild-type cells (IP shown in blue; Input shown in grey) and calibrated with *S. pombe* Scc1 distribution in representative chromosome V (IP shown in blue, bottom panel)

6

Measure DNA concentration using a Qubit HS assay kit.