



Version 2 ▾

Sep 09, 2022

Criteria for performing total knee arthroplasty without postoperative intensive care monitoring V.2

Fabricio Loures¹, João Henrique Reis¹, Marcelo de Oliveira Campos¹, Guilherme Morgado Runco¹, Luísa Doim Vieira Fonseca¹, Luciana Ferreira Xavier¹, Nelson Hiroyuki Miyabe Ooka¹, Liszt Palmeira de Oliveira¹

¹Universidade do Estado do Rio de Janeiro (UERJ)

1 Works for me

Share

dx.doi.org/10.17504/protocols.io.bp2l6b325gqe/v2



Fabricio Loures

Universidade do Estado do Rio de Janeiro (UERJ), Hospital Sa...

ABSTRACT

After reviewing the medical literature, we defined objective criteria to indicate intensive care monitoring after total knee arthroplasty. These criteria were adopted in January 2020 and since then none of the 80 patients who underwent total knee arthroplasty at the Pedro Ernesto University Hospital and who met the criteria required intensive care monitoring.

ATTACHMENTS

[Criterias for TKA without ICM.pdf](#)

DOI

dx.doi.org/10.17504/protocols.io.bp2l6b325gqe/v2

PROTOCOL CITATION

Fabricio Loures, João Henrique Reis, Marcelo de Oliveira Campos, Guilherme Morgado Runco, Luísa Doim Vieira Fonseca, Luciana Ferreira Xavier, Nelson Hiroyuki Miyabe Ooka, Liszt Palmeira de Oliveira 2022. Criteria for performing total knee arthroplasty without postoperative intensive care monitoring.

protocols.io

<https://protocols.io/view/criteria-for-performing-total-knee-arthroplasty-wi-cgdts6n>

Version created by Fabricio Loures



KEYWORDS

total knee arthroplasty, intensive care monitoring, risk stratification

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Sep 09, 2022

LAST MODIFIED

Sep 09, 2022

PROTOCOL INTEGER ID

69779

GUIDELINES

To perform total knee arthroplasty without postoperative intensive care monitoring, it is necessary for the patient to fulfill the eight criteria defined below:

Less than 75 years old

American Society Anesthesiology (ASA) Classification I or II

Non-smoker

No history of ischemic disease (Acute Myocardial Infarction and/or Cerebrovascular disease)

Primary Arthroplasty (CR or PS)

Body Mass Index (BMI) less than 35 kg/m²

Creatinine clearance higher than 60 ml/min

Hemoglobin greater than 12g/dL

BEFORE START

We have defined these criteria based on medical literature and we are validating this protocol by observing the outcome.

SAFETY WARNINGS

We have defined these criteria based on medical literature and we are validating this protocol by observing the outcome.