


Apr 26, 2024

Crystallization of Zika NS5 RdRp

DOI

dx.doi.org/10.17504/protocols.io.n2bvjnxjngk5/v1



anu.chandran^{1,2}, Peter Marples^{2,1}, Martin Walsh^{2,1}

¹Research Complex at Harwell; ²Diamond Light Source

anu.chandran: The principle crystallographer for the Zika NS5 RdRp polymerase project.;

ASAP Discovery



Lizbé Koekemoer

University of Oxford

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.n2bvjnxjngk5/v1

External link: <https://asapdiscovery.org/outputs/target-enabling-packages/#ASAP-ZIKA-NS5-RDRP>

Protocol Citation: anu.chandran, Peter Marples, Martin Walsh 2024. Crystallization of Zika NS5 RdRp. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.n2bvjnxjngk5/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: April 26, 2024

Last Modified: April 26, 2024

Protocol Integer ID: 98839

Keywords: crystallisation, XChem, ASAP, AViDD, CMD, Diamond Light Source, i04-1, Zika NS5 RdRp, NS5 RdRp, Research complex at Harwell

Funders Acknowledgement:

National Institutes of Health/National Institute Of Allergy and Infectious Diseases (NIH/NIAID)

Grant ID: Grant ID:
U19AI171399

Disclaimer

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Acknowledgements:

Diamond Light Source Ltd, Harwell Science and Innovation Campus, Didcot OX11 0QX, UK
Research Complex at Harwell, Harwell Science and Innovation Campus, Didcot OX11 0FA, UK
Oxford Lab Technologies crystal shifter <https://doi.org/10.1107/S2059798320014114>


Abstract

The main aim of this work was to identify small molecules that bind Zika NS5 RdRp (catalytic RNA-dependent RNA polymerase domain) through X-ray fragment-based screening. The Zika NS5 RDRP domain was cloned, expressed, purified, and crystallised. Suitable crystals for fragment screening were produced and optimised allowing an extensive fragment campaign to be performed. A native high-resolution structure was determined at 1.8Å and formed the basis for the fragment campaign.

Materials


SwissCI 3 lens crystallization plates <https://swissci.com/product/3-lens-crystallisation-plate/> **Codes:**
Midi: UVXPO-3LENS 3W96T-PS 3W96T-UVP

Morpheus HT-96 single reagent 250mL Catalog # MDSR-47-250-2-10

Purified Zika NS5 polymerase protein ([M] 5 mg/mL) in [M] 20 millimolar (mM) HEPES  7.5 ,
[M] 300 millimolar (mM) NaCl, 2.5% Glycerol, [M] 10 micromolar (µM) ZnCl₂, [M] 2 millimolar (mM) TCEP

306-903 residues- construct 2A1 (6 Hist sumo tag)

Safety warnings

 Follow all handling warning for the chemicals used in the crystallisation screen composition.

Equipment needed

- 1 **Formulatrix Rock Imager** (or incubator of choice)
SPT mosquito

Equipment

Mosquito HV

NAME

High Volume 16-Channel Robotic Liquid Handler

TYPE

SPT LabTech

BRAND

3097-01057

SKU

<https://www.sptlabtech.com/products/liquid-handling/mosquito-hv/>^{LINK}




P100 8 multi-channel pipette

SwissCI 3 lens plate

Crystallization experiment

1d


- 2 **Protein and buffer requirements:**



 28.8 µL  5 mg/mL  Sample

 3.264 mL Crystallization screen

- 3 **Crystallisation screen composition:**

Morpheus I E10 condition

 0.12 Molarity (M) Ethylene glycols

 0.1 Molarity (M) Buffer system 3  8.5

30 % v/v Precipitant Mix 2


Stock solutions used:




Morpheus I E10 condition

**Note**

The crystallisation screen can be stored in a duran bottle.

For long term storage keep the Crystallisation screen in the fridge at 4°C.

4 Dispense  34 µL Crystallisation screen into SwissCI 3 lens plate reservoir wells using a 100 µl multi-channel pipette.

Dispense  100 undetermined  5 mg/mL  Sample to each lens using the SPT mosquito.

Dispense  50 undetermined Crystallisation screen to each lens using the SPT mosquito.

Drop ratio: 2:1 ratio (100 nl  Sample : 50 nl reservoir solution)

Final drop volume: 150 nl

5 Incubate at  20 °C for  24:00:00 h in Formulatrix Rock Imager.

1d

Imaging Schedule: The first images are taken after 12 h and the imaging schedule follows a Fibonacci sequence of days for further collections.

6 Crystal form after ~24 h.

Expected result

The crystals reach their maximum size after 96 h and the precipitant has gone.

Crystals grown inconsistently to 2 sizes, one being half the size described below, but both sizes achieve the same results

Morphology: typically plates.


Size: ~ 250 μm in length and ~60 μm in width, depth of the crystals is ~2 μm

Appearance: glass shard.

Average resolution: 2.0 \AA

Space group: $P4_32_12$

Unit cell: 79 \AA , 79 \AA , 210 \AA
90.00°, 90.00°, 90.00°



An example of a drop containing Zika NS5 RdRp polymerase crystals.



Data collection at Synchrotron

7 Diamond Light Source
Unattended Data Collection (UDC)
Data Collection Temperature: 100K
Detector: DECTRIS EIGER2 X 9M
Beamline: I04-1
Wavelength: 0.9212 Å
Resolution (Å): 1.78
Beam Size (µm): 60 X 50
Number of images: 3600
Oscillation: 0.10°
Exposure (s): 0.0020
Transmission (%): 100
Flux (ph/s): 9.50e+11

Protocol references

N/A