

Apr 25, 2024

KAPP-Sen TMC: Term Placental Tissue Collection

DOI

dx.doi.org/10.17504/protocols.io.bp2l62b9dgqe/v1

Sonja Suvakov¹, Elizabeth Ann L. Enninga¹, Bridget A. Schley¹, Janelle Santos¹, Vesna D. Garovic¹

¹Mayo Clinic, Rochester, Minnesota, USA

Cellular Senescence Net...

KAPP-Sen TMC



Sergii Domanskyi

The Jackson Laboratory for Genomic Medicine

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.bp2l62b9dgqe/v1**

Protocol Citation: Sonja Suvakov, Elizabeth Ann L. Enninga, Bridget A. Schley, Janelle Santos, Vesna D. Garovic 2024. KAPP-Sen TMC: Term Placental Tissue Collection. **protocols.io** **<https://dx.doi.org/10.17504/protocols.io.bp2l62b9dgqe/v1>**

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: April 22, 2024

Last Modified: April 25, 2024

Protocol Integer ID: 98624

Keywords: SenNet KAPP-Sen TMC

Funders Acknowledgement:

National Institute on Aging

(NIA) KAPP-Sen Tissue

Mapping Collaborative


Grant ID: U54 AG075941



Abstract

The placenta samples are obtained within 15 minutes after delivery. As soon as the placenta is delivered it is placed in the box at room temperature and brought to the room where the samples are collected and labelled. Approximate time needed to complete the collection is 30 minutes.

1



Approximate site of collection.

- 2 Maternal side samples were collected within 3-5 cm range from the place where UC would be projected, cut into ~1 cm cubes and placed either in the cold MACS or cassettes with 10% buffered formalin.
- 3 Dissection of the fetal side of the placenta were made within 3-5 cm range from the UC and cut into ~1 cm cubes and placed either in the cold MACS or cassettes with 10% buffered formalin.
- 4 For the umbilical cord we cut the part that was closest to the baby. A total of ~3cm was cut and divided into the smaller pieces. The samples were immediately stored either in the cold MACS or cassettes with 10% buffered formalin.
- 5 Chorion was sampled from the fetal side per the instructions of Dr. Santos. It may contain the pieces of amniotic membrane as it was difficult to remove one from the other. A 3 cm cube was removed and divided into three smaller pieces which were stored as described in the previous steps. Note: fetal side=chorion



- 6 For the amniotic membrane we collected the very distant section that was wrapped around baby (edge). Same storage/transfer to the lab.
- 7 Upon arriving to the lab, all samples were prepared for shipping. Tissue stored in the cold MACS was transferred to new tubes wither with fresh, cold MACS (for scRNAseq) or flash frozen (for snRNAseq). Cassettes initially placed in the formalin jar were transferred to new formalin jars.