

Jul 05, 2024

## Immunohistochemistry for brain sections

DOI

**dx.doi.org/10.17504/protocols.io.3byl4956jgo5/v1**

Santiago Unda<sup>1,2</sup>, Roberta Marongiu<sup>1,2</sup>, Michael G. Kaplitt<sup>1,2</sup>

<sup>1</sup>Department of Neurosurgery, Weill Cornell Medical College, New York, NY 10065;

<sup>2</sup>Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD 20815, USA

ASAP Collaborative Rese...

Kaplitt Protocols



Eileen Ruth Torres

Weill Cornell Medicine

OPEN  ACCESS



DOI: [dx.doi.org/10.17504/protocols.io.3byl4956jgo5/v1](https://dx.doi.org/10.17504/protocols.io.3byl4956jgo5/v1)

**Protocol Citation:** Santiago Unda, Roberta Marongiu, Michael G. Kaplitt 2024. Immunohistochemistry for brain sections. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.3byl4956jgo5/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working

**We use this protocol and it's working**

**Created:** July 05, 2024

**Last Modified:** July 05, 2024

**Protocol Integer ID:** 102917

**Keywords:** ASAPCRN, immunohistochemistry, brain, mouse

**Funders Acknowledgement:**

**Aligning Science Across**

**Parkinson's**

**Grant ID:** 020608



## Abstract

Protocol for processing 30um mouse brain sections for immunolabeling.

## Materials

### **Solutions:**

#### **TBS-T**

For **2l**:

50mM Tris pH 7.5 – 100 ml of 1M stock

150 mM NaCl – 17.53 g

Triton 0.2 % - 4 ml

DW – 1.9 l

#### **ANTI-FREEZE SOLUTION**

For **1l**:

400 ml PBS/TBS (no TritonX-100!)

300 ml ethylene glycol



300 ml glycerine

STIR SOLUTION!!

#### **Blocking solution**

Prepare the solution in falcon tubes:

10 % goat, donkey or horse serum in TBS-T

Attn.: Do not shake too much (mix by inverting only). Proteins generate bubbles. Between experiments keep the blocking solution in the refrigerator at  4 °C . During experiment keep  On ice .



## Tissue Preparation

- 1 **Sections at cryostat:**
  - Thickness 30 µm
  - Sections picked up with a wet brush (not too wet) and moved to the TBS-T wells (usually serial sections). Store section in anti-freeze solution in -20 °C if not immediately staining.
- 2 **Transfer the sections** into the net-wells.

## Day 1: Primary Antibody Incubation

- 3 **Washes:** 3-step wash, 00:10:00 each, on a shaker, in TBS-T. (Between washes, just move the inserts from one tray to another.) 10m
- 4 **Blocking step:** add blocking solution into a 6-well plate.  
Transfer the sections with a clean brush from the net-inserts into the 6-well plate.  
Incubate on the shaker for 01:00:00 at Room temperature . 1h
- 5 **Primary antibody:** Transfer the sections into a 6-well plate containing the primary antibody solution (1 ml per well).
- 6 **Incubation:** on a shaker **in the cold room** Overnight . 1h

## Day 2: Secondary Antibody Incubation

- 7 **Washes:** a 3-step wash, 00:10:00 each, in TBS-T. 10m
- 8 **Secondary antibody:** Transfer the sections into a 12-well plate containing the secondary antibody. The secondary antibody will be incubated at Room temperature .
- 9 **Incubation:** on a shaker, at Room temperature , for 1-2 hours.



- 10 **Washes:** 3-4 times, for ⌚ 00:10:00 , in TBS-T. 10m
- 11 **DAPI:** incubate with DAPI (1:10,000) shaking for ⌚ 00:10:00 at 🌡 Room temperature . 10m
- 12 **Washes:** a 2-step wash, on a shaker, at 🌡 Room temperature , ⌚ 00:10:00 each. 10m
- 13 **Mounting:** Mount sections and let them dry for ⌚ 00:10:00 at 🌡 Room temperature . 10m
- 14 **Coverslip:** Let the slides dry at least ⌚ Overnight at 🌡 Room temperature . 10m