

Jul 11, 2024

Mouse brain slice electrophysiology

DOI

dx.doi.org/10.17504/protocols.io.rm7vzj7y8lx1/v1

Shiyi Wang¹

¹Duke University

ASAP Collaborative Rese...



Shiyi Wang

Duke University

OPEN  ACCESS



DOI: dx.doi.org/10.17504/protocols.io.rm7vzj7y8lx1/v1

Protocol Citation: Shiyi Wang 2024. Mouse brain slice electrophysiology. **protocols.io**

<https://dx.doi.org/10.17504/protocols.io.rm7vzj7y8lx1/v1>

License: This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: July 11, 2024

Last Modified: July 11, 2024

Protocol Integer ID: 103202

Keywords: ASAPCRN

Funders Acknowledgement:
Aligning Science Across
Parkinson's (ASAP) initiative
Grant ID: ASAP-020607



Disclaimer

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to **protocols.io** is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with **protocols.io**, can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

Abstract

Mouse brain slice electrophysiology

1 ****Animal Handling****

- 1.1 - Anesthetize 3-4 mice of each genotype (WT and LRRK2 G2019Ski/ki) and condition using 200 mg/kg tribromoethanol (Avertin).
- 1.2 - Ensure proper anesthesia depth and confirm absence of reflexes before proceeding.

2 ****Brain Extraction and Slicing****

- 2.1 - Decapitate the mice and immediately immerse the brains in ice-cold artificial cerebrospinal fluid (aCSF) containing (in mM): 125 NaCl, 2.5 KCl, 3 mM MgCl₂, 0.1 mM CaCl₂, 10 glucose, 25 NaHCO₃, 1.25 NaHPO₄, 0.4 L-ascorbic acid, and 2 Na-pyruvate, pH 7.3-7.4 (310 mOsmol).
- 2.2 - Use a vibrating tissue slicer (Leica VT1200) to obtain 350 µm thick coronal brain slices containing the anterior cingulate cortex (ACC).

3 ****Slice Incubation****

- 3.1 - Transfer slices immediately to standard aCSF maintained at 33°C and continuously bubbled with 95% O₂ – 5% CO₂.
- 3.2 - Incubate slices for 30 minutes at 33°C to recover.

4 ****Transfer to Holding Chamber****

- 4.1 - After incubation, transfer slices to a holding chamber at room temperature (approximately 25°C) with the same extracellular buffer.

5 ****Microscopy Setup****

- 5.1 - Visualize brain slices using an upright microscope (BX61WI, Olympus) equipped with a 40x water-immersion objective and infrared-differential interference contrast optics.

5.2 - Use a digital camera (ODA-IR2000WCTRL) for image acquisition.

6 ****Patch-Clamp Setup****

6.1 - Perform patch-clamp recordings using an EPC 10 patch-clamp amplifier controlled by Patchmaster Software (HEKA).

6.2 - Set data acquisition at a sampling rate of 50 kHz and low-pass filter at 6 kHz.

7 ****Recording Configuration****

7.1 - Use aCSF bath solution containing: For mEPSCs: 1 μ M tetrodotoxin and 50 μ M Picrotoxin, held at -70 mV in voltage-clamp mode; For mIPSCs: 1 μ M tetrodotoxin, 10 μ M CNQX, and 50 μ M D-AP5, held at -70 mV in voltage-clamp mode.

8 ****Internal Solution Preparation****

8.1 - Prepare internal solutions for patch pipettes: For mEPSCs: 125 K-gluconate, 10 NaCl, 10 HEPES, 0.2 EGTA, 4.5 MgATP, 0.3 NaGTP, and 10 Na-phosphocreatine, pH adjusted to 7.2 – 7.4 with KOH, osmolality ~300 mOsmol. For mIPSCs: 77 K-gluconate, 77 KCl, 10 HEPES, 0.2 EGTA, 4.5 MgATP, 0.3 NaGTP, and 10 Na-phosphocreatine, pH adjusted to 7.2 – 7.4 with KOH, osmolality ~300 mOsmol.

9 ****Data Acquisition and Analysis****

9.1 - Record mEPSCs and mIPSCs using Minhee Analysis software.

9.2 - Analyze frequency by counting events over 5 minutes of recording.

9.3 - Calculate average events per cell based on at least 100 non-overlapping events.



9.4 - Measure peak amplitude of average mEPSCs relative to baseline current.