



Mar 06,  
2020

## Extraction Procedure

Jing Xu<sup>1</sup>

<sup>1</sup>Hainan University

1

Works for me

[dx.doi.org/10.17504/protocols.io.bdawi2fe](https://doi.org/10.17504/protocols.io.bdawi2fe)



Jing Xu  
Hainan University



### ABSTRACT

Dried leaves of *C. cyrtophyllum* (150 g) were weighted and sieved (20 mesh) in an herb disintegrator (118 Swing, Zhejiang, China); the powdered samples were extracted twice according to previous protocol [10]. Solvent was removed from the combined filtrates and 61.44 g of ECE was obtained and redissolved in distilled water (500 ml). The solution was partitioned with 3×250 ml petroleum ether (60–90 °C), 3×250 ml dichloromethane, 3×250 ml ethyl acetate and 3×250 ml *n*-butanol. The resulting extracts were concentrated to yield sub-fractions, PEF, DMF, EAF, BAF and RF, which were 0.65, 5.53, 4.13, 13.28 and 36.97 g, respectively. The samples were stored at 4 °C.



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited