



Mar 13, 2021

Streptococcal Protein-G and Protein-LG sandwich ELISA

Angel A Justiz-Vaillant¹¹University of the West Indies St. Augustine

1

Works for me

dx.doi.org/10.17504/protocols.io.btaznif6

Carbon

Angel Justiz-Vaillant

University of the West Indies St. Augustine

SUBMIT TO PLOS ONE

ABSTRACT

This ELISA was used to study the interactions between SpG) and SpLG) with different immunoglobulin preparations from mammalian and avian species.

DOI

dx.doi.org/10.17504/protocols.io.btaznif6

PROTOCOL CITATION

Angel A Justiz-Vaillant 2021. Streptococcal Protein-G and Protein-LG sandwich ELISA. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.btaznif6>

LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Mar 13, 2021

LAST MODIFIED

Mar 13, 2021

PROTOCOL INTEGER ID

48185

- 1 This ELISA was used to study the interactions between SpG) and SpLG) with different immunoglobulin preparations from mammalian and avian species. The 96 well microtiter plate was coated overnight at 4°C with 2 µg/µl per well of SpG in carbonate-bicarbonate buffer pH 9.6.
- 2 The plate was then treated with bovine serum albumin solution and washed 4X with PBS-Tween. 50 µl of immunoglobulins (1 mg/ml) is added and incubated for 1h at room temperature, and the microplate is rewashed 4X with PBS-Tween.
- 3 Then, 50 µL of peroxidase-labeled SpLG conjugate diluted 1:5000 in PBS-non-fat milk was added to each well and incubated for 1h at RT. The plate was washed 4X with PBS-Tween.

- 4 Then, 50 μ L of o-phenylenediamine solution (4 mg/mL) was added, and the plate was incubated for 15 min at RT in the dark. The reaction was stopped with 50 μ L of a 3M H₂SO₄ solution.
- 5 The plate was visually assessed for color development and read on a microplate reader at 492 nm. A cut-off point was calculated as the mean of the optical density of the negative controls multiplied by two. The cut-off value was set to 0.36.