

AUG 10, 2023

OPEN BACCESS



DOI:

dx.doi.org/10.17504/protocol s.io.bp2l6x955lqe/v1

Protocol Citation: Andriko Palmowski, Judith Oldenkott, Anne Pankow, Anne Beenken, Eric L Matteson, Frank Buttgereit 2023. Confounding by Indication in Observational Studies Investigating Glucocorticoid-Associated Adverse Events in Rheumatoid Arthritis: A Protocol for a Systematic Review. protocols.io https://dx.doi.org/10.17504/protocols.io.bp2l6x955lqe/v1

License: This is an open access protocol distributed under the terms of the Creative Commons
Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

© Confounding by Indication in Observational Studies Investigating Glucocorticoid-Associated Adverse Events in Rheumatoid Arthritis: A Protocol for a Systematic Review

Andriko Judith

Palmowski¹, Oldenkott¹, Anne Pankow¹, Anne Beenken¹, Eric L Matteson², Frank Buttgereit¹

¹Department of Rheumatology and Clinical Immunology, Charité - University Medicine Berlin, Germany;

²Mayo Clinic, Rochester, Minnesota, United States of America



Andriko Palmowski

Department of Rheumatology and Clinical Immunology, Charité ...

ABSTRACT

_

Protocol status: Working We use this protocol and it's

working

Created: Aug 10, 2023

Last Modified: Aug 10,

2023

PROTOCOL integer ID:

86324

Abstract

Background Glucocorticoids (GCs) are regularly used drugs in rheumatoid arthritis (RA), but they can cause adverse events (AEs). Observational studies provide important real-world evidence but can be biased. Observational studies investigating the association between AEs and GCs in RA are

particularly susceptible to confounding by indication as patients with higher disease activity are more likely to suffer from AEs and more likely to receive GCs.

Objective To assess how often observational studies in RA which investigate the association between GCs and certain AEs adjust for disease activity (such as the disease activity score-28 joints) and/or systemic inflammation (such as serum C-reactive protein levels).

Methods A systemic literature review will be conducted and reported in compliance with the 2020 PRISMA statement.

PDF Protocol

2

Protocol.pdf