



SEP 26, 2023

OPEN ACCESS



**DOI:**  
[dx.doi.org/10.17504/protocols.io.5qpvo3o7xv4o/v1](https://dx.doi.org/10.17504/protocols.io.5qpvo3o7xv4o/v1)

**Protocol Citation:** NUS iGEM 2023. Preparation of Agarose Gel. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.5qpvo3o7xv4o/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

**Protocol status:** Working  
 We use this protocol and it's working

**Created:** Sep 26, 2023

**Last Modified:** Sep 26, 2023

## 🌐 Preparation of Agarose Gel

NUS iGEM<sup>1</sup>

<sup>1</sup>National University of Singapore

NUS iGEM 2023



NUS iGEM

National University of Singapore

### DISCLAIMER

DISCLAIMER – FOR INFORMATIONAL PURPOSES ONLY; USE AT YOUR OWN RISK

The protocol content here is for informational purposes only and does not constitute legal, medical, clinical, or safety advice, or otherwise; content added to [protocols.io](#) is not peer reviewed and may not have undergone a formal approval of any kind. Information presented in this protocol should not substitute for independent professional judgment, advice, diagnosis, or treatment. Any action you take or refrain from taking using or relying upon the information presented here is strictly at your own risk. You agree that neither the Company nor any of the authors, contributors, administrators, or anyone else associated with [protocols.io](#), can be held responsible for your use of the information contained in or linked to this protocol or any of our Sites/Apps and Services.

### ABSTRACT

2023 NUS-Singapore iGEM team followed this protocol to prepare a 1% agarose gel. The team used this 1% agarose gel to isolate the DNA fragments from the common PCR and the colony PCR products.




### MATERIALS

1. Agarose Powder
2. 1x TAE Buffer (Tris-Acetate-EDTA Buffer)
3. Ultra GelRed Gel Stain


**Keywords:** Agarose Gel, Gel, Agarose, Gel Electrophoresis, DNA Isolation



Proper lab PPE must be worn at all times while preparing the agarose gel. Additionally, thermal gloves designed for high-temperature protection must be worn when handling the hot agarose solution after microwaving.

- 1 For a 1% agarose gel, mix the following in a conical flask:
    -  5 g of Agarose powder.
    -  50 mL of 1x TAE buffer (Tris-Acetate-EDTA Buffer).
  - 2 Swirl the agarose solution to mix it well.
  - 3 Heat the agarose solution in a microwave until it boils.
  - 4 Take out the conical flask from the microwave and swirl the conical flask until the agarose solution is clear and without undissolved agarose powder or lumps.
- Safety information**
- Wear thermal gloves when handling the hot conical flask from the microwave to prevent burns.
- 5 Add  5  $\mu$ L of Ultra GelRed gel stain into the agarose solution and swirl the conical flask until the colour becomes uniform.
  - 6 Secure a gel tray tightly onto a gel caster and place an 8-well comb or 15-well comb into the tray.

**7** Pour the agarose solution into the tray, ensuring that there are no bubbles.

**8** Cool down the agarose solution for at least  00:30:00 to get a solidified agarose gel.

30m