

Sep 30, 2024

Genomic DNA Extraction from Sorted Cells

 In 1 collection

DOI

dx.doi.org/10.17504/protocols.io.n2bvjn88wgk5/v1

Raining Wang¹, Melinda Wheelock¹

¹University of Washington, Department of Genome Sciences

IGVF



Sarah Heidl

University of Washington, Starita Lab

OPEN  ACCESS



DOI: **dx.doi.org/10.17504/protocols.io.n2bvjn88wgk5/v1**

Protocol Citation: Raining Wang, Melinda Wheelock 2024. Genomic DNA Extraction from Sorted Cells. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.n2bvjn88wgk5/v1>

License: This is an open access protocol distributed under the terms of the **[Creative Commons Attribution License](#)**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: May 09, 2024

Last Modified: September 30, 2024

Protocol Integer ID: 99787

Funders Acknowledgement:

NHGRI (IGVF)

Grant ID: HG011969



Abstract



This protocol details the extraction of genomic DNA from cells.

Guidelines

The DNeasy columns cannot take more than 5×10^6 cells per column. Larger pellets can be split into multiple extraction reactions, then combined after elution.

Materials

Materials

- Pelleted cells for DNA extraction (thawed, if previously frozen)
-  DNeasy Blood & Tissue Kits **Qiagen Catalog #69506**
- 1.7-mL microcentrifuge tubes
- DPBS
-  Monarch RNase A – 1 ml (2x0.5ml) **New England Biolabs Catalog #T3018L**
- Molecular grade water
- 2.0-mL microcentrifuge tubes
- 96-100% Ethanol

Equipment

- Benchtop centrifuge
- Thermomixer
- Vortexer
- Picofuge
- Nanodrop




Extraction of genomic DNA from cells

39m


- 1 Prepare a lysis master mix of the following reagents and mix by vortexing briefly:




A	B	C
Reagent	Volume per sample (uL)	Volume for X samples (uL)
DPBS	100	$X * 100 * 1.05$
Buffer AL	200	$X * 200 * 1.05$
Proteinase K	20	$X * 20 * 1.05$
RNase A	2	$X * 2 * 1.05$
Total	322	



- 2 Aliquot  322 μL of this master mix into labeled 1.7-mL microcentrifuge tubes.

- 3 Resuspend the pellets in  100 μL DPBS for every 5×10^6 cells in each sample pellet.

- If you have 10×10^6 cells in your pellet, resuspend in  200 μL , then split into 2 separate tubes.


- 4 Add resuspended cells in  100 μL DPBS to an appropriately labeled tube containing the lysis master mix. Mix by vortexing.



- 5 Incubate the resuspended cells at  56 °C with agitation at  1500 rpm, 00:30:00 .

30m



- 6 Spin down the lysis tubes briefly, then add  200 μL of 96-100% ethanol to the lysed cells. Mix thoroughly by vortexing. Spin the tubes down to remove drops from lids.














- 7 Transfer the lysed cell mixture to a labeled DNeasy mini spin column in a 2mL collection tube (both provided in the kit).

- 8 Centrifuge at  6000 x g, 00:01:00 . Discard the flow-through and collection tube.

1m



- 9 Place the spin column in a new 2mL collection tube and add  500 μ L Buffer AW1 (provided in kit) to the column. Centrifuge at  6000 x g, 00:01:00 . Discard the flow-through and collection tube.
- 10 Place the spin column in a new 2 mL collection tube and add  500 μ L Buffer AW2 (provided in kit) to the column. Centrifuge at  20000 x g, 00:03:00 . Discard the flow-through and collection tube.
- 11 Place the spin column in a new 2mL collection tube and centrifuge at  20000 x g, 00:01:00 .
- 12 Place the spin column in a new (labeled) 1.7-mL microcentrifuge tube.
- 13 Add  100 μ L of molecular grade water to the spin column and incubate at  Room temperature for  00:02:00 . Centrifuge at  6000 x g, 00:01:00 .
- 14 Repeat Step 13 with an additional  100 μ L of molecular grade water.
- 15 Transfer any split samples (pellets with more than 5×10^6 cells) into labeled 2.0mL microcentrifuge tubes.
- 16 Quantitate with the Nanodrop, and store the samples at  -20 $^{\circ}$ C .



1m



3m



1m



3m

