



Oct 13, 2020

DNA-Extraction-ClostridiaProject

Tobias Wandt¹¹Universität Leipzig

1

Works for me

This protocol is published without a DOI.



Tobias Wandt

ABSTRACT

Protocol used for DNA-Extraction of Clostridia in my project.

DNA-Kit used is GenFind V3 by Beckmann/Coulter.

Some steps used from:

Sim JH, Anikst V, Lohith A, Pourmand N, Banaei N. Optimized Protocol for Simple Extraction of High-Quality Genomic DNA from Clostridium difficile for Whole-Genome Sequencing. J Clin Microbiol. 2015;53(7):2329-2331. doi:10.1128/JCM.00956-15

PROTOCOL CITATION

Tobias Wandt 2020. DNA-Extraction-ClostridiaProject. **protocols.io**
<https://protocols.io/view/dna-extraction-clostridiaproject-bnb9mar6>



LICENSE

————— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 12, 2020

LAST MODIFIED

Oct 13, 2020

PROTOCOL INTEGER ID

43105

ABSTRACT

Protocol used for DNA-Extraction of Clostridia in my project.

DNA-Kit used is GenFind V3 by Beckmann/Coulter.

Some steps used from:

Sim JH, Anikst V, Lohith A, Pourmand N, Banaei N. Optimized Protocol for Simple Extraction of High-Quality Genomic DNA from Clostridium difficile for Whole-Genome Sequencing. J Clin Microbiol. 2015;53(7):2329-2331. doi:10.1128/JCM.00956-15

Growing and Preparing Clostridia

2d

2d

- 1 Inoculate Clostridia in 10ml TSB-Broth, incubating at least 48h in anaerobe atmosphere at 37°C

10 mL TSB broth **48:00:00 at 37°C anaerobic**

15m

- 2 Take 1ml of Clostridia TSB-Broth and centrifuge for 15min at 3,000g, after that remove supernatant

1 mL Clostridia TSB broth **00:15:00 centrifuge at 3000g**

- 3 Add 500ul Lysis-Buffer(LBB) and 30ul Proteinase K, tipmixing or vortexing to resuspend the pellet
 **500 µl LBB**  **30 µl Proteinase K**
- 4 Incubate at 57°C for 1h while continuously shaking at ~500/min 1h
 **01:00:00 at 57°C + shaking 500/min**
- 5 Add 300ul vortexed Black Beads(BBB) and tipmix(at least 10 times), incubate for 5 min at room temperature, place on a magnetic tube stand until no or few beads are left in suspension ~10-15min, then carefully remove supernatant 20m
without disturbing the beads-formation
 **300 µl Black Beads**  **00:05:00 at room temperature**  **00:15:00 magnetic tube stand**
- 6 Add 800ul WBB and tipmix(at least 10 times), place on a magnetic tube stand until no or few beads are left in suspension ~5min, remove supernatant 5m
 **800 µl WBB**  **00:05:00 magnetic tube stand**
- 7  **go to step #6 for a total of 2 WBB washes**
- 8 Add 800ul WBC and tipmix (at least 10 times), than add 600ul WBC and tipmix, place on a magnetic tube stand until no or few beads left in suspension ~5min , remove supernatant 5m
 **1400 µl WBC**  **00:05:00 magnetic tube stand**
- 9  **go to step #8 for a total of 2 WBC washes**
- 10 Add 100ul distilled water, tipmix 10 times, incubate for 2min at roomtemp, tipmix 10 times again and place in a 4°C 2m
fridge overnight
 **100 µl distilled water**  **00:02:00 room temperature**  **Overnight at 4°C**
- 11 Tipmix 10times and place on a magnetic tube stand until no beads are left in suspension
- 12 Take the supernatant with your DNA and put it in a fresh tube