



Jun 03, 2020

Degassing Reagents

Allen Institute for Brain Science¹

¹Allen Institute

1 Works for me

dx.doi.org/10.17504/protocols.io.bg4wjyxe

BICCN Allen Institute for Brain Science



ABSTRACT

This protocol is used for the degassing of critical reagents used in both the colorimetric and fluorescent in situ hybridization (ISH). This is crucial in eliminating hybridization bubbles from forming on the tissue. Hybridization bubbles result in no expression where they are located on the tissue.

Note: Research reported in this publication was supported by the National Institute Of Mental Health of the National Institutes of Health under Award Number U19MH114830. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

ATTACHMENTS

RP0017_Degassing_Reage nts.docx

DOI

dx.doi.org/10.17504/protocols.io.bg4wjyxe

PROTOCOL CITATION

Allen Institute for Brain Science 2020. Degassing Reagents. **protocols.io** dx.doi.org/10.17504/protocols.io.bg4wjyxe

KEYWORDS

RP0017, in situ hybridization, ISH

LICENSE

This is an open access protocol distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Jun 03, 2020

LAST MODIFIED

Jun 03, 2020

PROTOCOL INTEGER ID

37750