



Oct 24, 2020

Gene knockout

Zhujun Wei¹¹2020 iGEM NEFU China

1

Works for me

dx.doi.org/10.17504/protocols.io.bnwzmff6

2020 iGEM NEFU China



Zhujun Wei

ABSTRACT

We use this protocol to knock out yhaK in E.coli BW25113

DOI

dx.doi.org/10.17504/protocols.io.bnwzmff6

PROTOCOL CITATION

Zhujun Wei 2020. Gene knockout. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bnwzmff6>

LICENSE

This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Oct 24, 2020

LAST MODIFIED

Oct 24, 2020

PROTOCOL INTEGER ID

43705

MATERIALS TEXT

E. coli BW25113
pCas9 plasmid
LB medium
Sterilized 10% glycerol
1mM IPTG
Kana
Str

SAFETY WARNINGS

We use this protocol to knock out yhaK in E.coli BW25113

ABSTRACT

We use this protocol to knock out yhaK in E.coli BW25113

BEFORE STARTING

Please read the protocol carefully before doing the experiment.

- 1 Transfer the pCas9 plasmid into the recipient E. coli BW25113 (Kana resistance, temperature sensitive).

2 Preparation of pCas9 competent cells:

- 2.1 Add 50 μ L of bacterial solution to 5 mL of LB medium, incubate at 30°C, 200 rpm for about 12 h, Add 2% to 100 mL LB medium, add 0.5% of arabinose at a final concentration after 0.5 hours of culture;
- 2.2 Incubate at 30°C for about 2~2.5 h, until the OD600 is about 0.55-0.6, then take it out and let it stand on ice for 30 min;
- 2.3 Collect cells at 4200 rpm for 10 min in a 4°C refrigerated centrifuge;
- 2.4 Resuspend the cells gently with 10 ml of pre-cooled and sterilized 10% glycerol on ice.
- 2.5 Repeat steps 2.3 and 2.4 three times, using 10 mL of pre-cooled and sterilized 10% glycerol for the first two washes. The last time it was resuspended with 0.5 mL of glycerol.
- 2.6 Dispense in 100 μ L/tube into a pre-cooled centrifuge tube, it can be used for electroporation or immediately placed in a -80°C refrigerator. The bacteria can be stored at -80°C for half a year.

3 Prepare target segment: Clone upstream and downstream sequences of yhaK by PCR to about 500 bp each, then connect upstream and downstream homology arms together by overlap.

4 pTarget-yhaK vector construction (Str resistance): Download the yhaK gene sequence, Find the appropriate gRNA affinity site and replace 1961 to 1980 bases of the pTarget plasmid by digestion and ligation or overlap.

5 Electroporation:

- 5.1 Add >200ng target fragment and >40ng pTarget-yhaK plasmid to 100 μ L competent cells on ice for 10~30 min, the plasmid and the ligation product should not exceed 5 μ L;
- 5.2 UV-sterilize the washed and dried 2 mm electroporation cuvette for 20 min and pre-cooled on ice, then quickly transfer the competent cells to the bottom of the electroporation cuvette;
- 5.3 Dry the outer wall of the cup. Immediately after the electric shock, add 900~1000 μ L of 37 °C preheated

LB medium, mix gently, transfer to a 1.5 mL centrifuge tube, shake at 30 °C, 150 rpm, 45 ~ 60 min, then apply an appropriate amount of bacterial solution to the plate containing Kana and Str.

- 6 Knockout verification: Primers were designed at 700-1000 bp upstream and downstream of the yhaK in the genome to perform colony PCR validation on clones on the plates.
- 7 pTarget-yhaK elimination: Take the correct clone, add 1mM IPTG at 30°C for 12 h, streak to isolate the monoclonal. The clones were tested for the presence of Str resistance, clones with pTarget eliminated is not resistant to str. The pCas9 plasmid can be eliminated by incubation at 37°C.
- 8 Strains that retain the pCas9 plasmid can be used to knock out other genes.