



SEP 19, 2023

OPEN  ACCESS**DOI:**

[dx.doi.org/10.17504/protocol  
s.io.bp2l6xdwrlqe/v1](https://dx.doi.org/10.17504/protocols.io.bp2l6xdwrlqe/v1)

**Collection Citation:** jlancieg  
o 2023. Protocol for  
"Neuromelanin accumulation  
drives endogenous  
synuclienopathy in non-  
human primates".

**protocols.io**

[https://dx.doi.org/10.17504/p  
rotocols.io.bp2l6xdwrlqe/v1](https://dx.doi.org/10.17504/protocols.io.bp2l6xdwrlqe/v1)

**License:** This is an open  
access collection distributed  
under the terms of  
the [Creative Commons  
Attribution License](#), which  
permits unrestricted use,  
distribution, and reproduction  
in any medium, provided the  
original author and source  
are credited

**Protocol status:** Working

We use this collection and it's  
working

**Created:** Sep 19, 2023


## Protocol for "Neuromelanin accumulation drives endogenous synuclienopathy in non-human primates"

jlanciego<sup>1,2,3</sup>

<sup>1</sup>CNS Gene Therapy Program, Center for Applied Medical Research (CIMA)  
University of Navarra;

<sup>2</sup>Centro de Investigación Biomédica en Red de Enfermedades  
Neurodegenerativas (CiberNed-ISCIII);

<sup>3</sup>Aligning Science Across Parkinson's (ASAP) Collaborative Research  
Network



jlanciego

**DISCLAIMER**

The authors report no competing interests

**Last Modified:** Sep 19, 2023

**COLLECTION integer ID:** 87996

**Keywords:** Tyrosinase, Alpha-synuclein, Lewy bodies, Marinesco bodies, Prion-like spread

## ABSTRACT

This study was aimed to develop and characterize a non-human primate (NHP) model of Parkinson's disease mimicking the known neuropathological hallmarks of Parkinson's disease to the best possible extent. Accordingly, we sought to determine whether AAV-mediated enhanced expression of human tyrosinase (hTyr) in the substantia nigra (SNpc) of non-human primates (NHPs) is able to induce a time-dependent accumulation of neuromelanin (NMel) in dopaminergic neurons, further triggering and endogenous synucleinopathy, progressive cell death and a pro-inflammatory scenario, in keeping with what was formerly reported in rats by taking advantage of a similar strategy (Carballo-Carbajal et al., 2019). Furthermore, the potential prionoid spread of endogenous alpha-synuclein (a-Syn) species towards the prefrontal cortex was analyzed, in an attempt to evaluate to what extent there is a propagation of endogenous a-Syn by permissive trans-synaptic templating (e.g. the so-called Braak hypothesis). Adult juvenile NHPs (*Macaca fascicularis*) were injected with adeno-associated viral vectors (AAVs) encoding either the hTyr gene (AAV-hTyr; delivered into the left SNpc) or a null construct for control purposes (AAV-null; injected into the right SNpc). In order to delineate a timeline for the underlying processes, one group of NHPs was sacrificed four months post-AAV deliveries (animals M308F4 and M310M4), whereby the follow-up timing for second experimental group was settled at eight months post-AAVs surgeries (animals M307F8 and M309M8). Neuroimage studies (MRI and MicroPET) were conducted *in vivo* at different time points. Upon animal sacrifices, brain tissue samples were processed for histological analysis comprising intracellular NMel levels, intracellular aggregates, nigrostriatal degeneration and neuroinflammation.

## ATTACHMENTS

[PROTOCOL.docx](#)

## DISCLAIMER

The authors report no competing interests

## ATTACHMENTS

PROTOCOL.d  
OCX

## FILES