



Oct 07, 2022

# PD-L1 expression as predictor of immunotherapy eligibility in penile squamous cell carcinoma patients

Muhammad Haritsyah Warli<sup>1</sup>, Fauriski Febrian Prapiska<sup>2</sup>,  
Ginanda Putra Siregar<sup>2</sup>, Bungaran Sihombing<sup>2</sup>, Syah Mirsyah Warli<sup>2</sup>

<sup>1</sup>Universitas Indonesia; <sup>2</sup>Universitas Sumatera Utara

Muhammad Haritsyah Warli: Department of Urology, Faculty of Medicine, Universitas Indonesia - Haji Adam Malik General Hospital;

Fauriski Febrian Prapiska: Division of Urology, Department of Surgery, Faculty of Medicine, Universitas Sumatera Utara – Haji Adam Malik General Hospital

Ginanda Putra Siregar: Division of Urology, Department of Surgery, Faculty of Medicine, Universitas Sumatera Utara – Haji Adam Malik General Hospital

Bungaran Sihombing: Division of Urology, Department of Surgery, Faculty of Medicine, Universitas Sumatera Utara – Haji Adam Malik General Hospital

Syah Mirsyah Warli: Division of Urology, Department of Surgery, Faculty of Medicine, Universitas Sumatera Utara – Haji Adam Malik General Hospital and Department of Urology, Faculty of Medicine, Universitas Sumatera Utara Hospital – Universitas Sumatera Utara

1 Works for me

Share

[dx.doi.org/10.17504/protocols.io.bp2l69b4klqe/v1](https://dx.doi.org/10.17504/protocols.io.bp2l69b4klqe/v1)

Muhammad Haritsyah Warli

## ABSTRACT

We investigated the correlation between PD-L1 expression and poor prognosis. We examined the correlation between PD-L1 expression and survival of penile SqCC and analyzed the adjusted hazard ratios (HR) of the patients with PD-L1 expression compared to anatomical stage of the tumor.

## DOI

[dx.doi.org/10.17504/protocols.io.bp2l69b4klqe/v1](https://dx.doi.org/10.17504/protocols.io.bp2l69b4klqe/v1)

## PROTOCOL CITATION

Muhammad Haritsyah Warli, Fauriski Febrian Prapiska, Ginanda Putra Siregar, Bungaran Sihombing, Syah Mirsyah Warli 2022. PD-L1 expression as predictor of immunotherapy eligibility in penile squamous cell carcinoma patients.

**protocols.io**

<https://dx.doi.org/10.17504/protocols.io.bp2l69b4klqe/v1>



## KEYWORDS

PD-L1, Squamous Cell Carcinoma

## LICENSE

\_\_\_\_\_ This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

## CREATED

Oct 05, 2022

## LAST MODIFIED

Oct 07, 2022

## PROTOCOL INTEGER ID

70872





## MATERIALS TEXT

- MD21R clone (Medaysis CA)
- Rabbit PD-L1 antibody
- DAB mixture
- Hematoxylin

### PD-L1 Antibody Preparation

- 1 Reagent used to stain PD-L1 in this IHC staining was MD21R clone (Medaysis CA), ready to use (rabbit PD-L1 antibody)
- 2 PD-L1 antibody is diluted with ratio 1 uL PD-L1: 100 uL IHC diluent and then mixed for 5 minutes. 5m

### Pretreatment

- 3 Pretreatment was done with EDTA pH8.0, 15 minutes using a Pressure Cooker 1h 15m  
  
 **00:15:00 Pressure cooker**  
OR  
  
 **01:00:00 30 – 60 minutes using a water bath**  **95 °C** to  **99 °C**
- 4 Deparaffination and rehydration were done by using Bond Dewax Solution for 3 times, alcohol for 3 times, and Bond Wash Solution for 3 times.

- 5 We retrieved the antigen by using Bond Epitope Retrieval Solution I for 4 times

#### PD-L1 and Sample Mixing and Detection

- 6 We mixed the solution with PD-L1 (Medaysis) primary antibody for 1 hour. 1h

- 7 The detection was done by using post primary procedure for 8 minutes, Bond Wash Solution<sup>16m</sup> for two times, and polymer for eight minutes. We used DAB mixture for staining and hematoxylin for counterstaining.

🕒 00:08:00 post primary procedure

Bond Wash Solution Two times

🕒 00:08:00 polymer

#### Analysis

- 8 PD-L1 expression was assessed on cytoplasm and/or tumor cell membranes and TILs. The positive control used was the placenta.
- 9 PD-L1 expression is considered positive if it has a score of +2 or +3 and is considered negative if it has a score of +1 and 0. Intensity measurement then given a score of 0-3 (0= none; 1= weak; 2= moderate; and 3= strong).
- 10 The percentage of stained cells was assessed on a scale of 0-100%.  
Percentage of stained tumor cells A= percentage of tumor cells intensity in 3+ {3 x (%Cells 3+)}, B= percentage of tumor cells intensity in 2+ {2 x (%Cells 2+)}, C= percentage of tumor cells intensity in 1+ {1 x (%Cells 1+)}
- 11 Semi-quantitative assessment refers to research by Chovanec *et al.* who used histoscore (H-score)  
$$\text{H-Score} = \{1 \times (\% \text{cells } 1+) + 2 \times (\% \text{cells } 2+) + 3 \times (\% \text{cells } 3+)\}$$
- 12 Total value obtained ranges from 0 to 300, thus PD-L1 expression is categorized into "1" for negative (0 – 99) and "2" for positive (100 – 300).

