




Nov 21, 2020

CELL COUNT- 02 - Manual cell count with Türk Solution

Marco Cosentino¹, Elisa Storelli¹, Alessandra Luini¹, Massimiliano LM Legnaro¹, Emanuela Rasini¹, Marco Ferrari¹, Franca Marino¹

¹Center for Research in Medical Pharmacology, University of Insubria (Varese)

1 Works for me dx.doi.org/10.17504/protocols.io.bpxsmpne

 Farmacologia Medica

ABSTRACT

Published work using this protocol:

- Kustrimovic N., Comi C., Magistrelli L., Rasini E., Legnaro M., Bombelli R., Aleksic I., Blandini F., Minafra B., Riboldazzi G., Struchio A., Mauri M., Bono G., Marino F., Cosentino M. Parkinson's disease patients have a complex phenotypic and functional Th1 bias: cross-sectional studies of CD4+ Th1/Th2/T17 and Treg in drug-naïve and drug-treated patients (2018). Journal of neuroinflammation, 15(1), 205. <https://doi.org/10.1186/s12974-018-1248-8>

- Kustrimovic, N., Rasini, E., Legnaro, M., Bombelli, R., Aleksic, I., Blandini, F., Comi, C., Mauri, M., Minafra, B., Riboldazzi, G., Sanchez-Guajardo, V., Marino, F., & Cosentino, M. (2016). Dopaminergic Receptors on CD4+ T Naive and Memory Lymphocytes Correlate with Motor Impairment in Patients with Parkinson's Disease. Scientific reports, 6, 33738. <https://doi.org/10.1038/srep33738>

- Cosentino M., Ferrari M., Kustrimovic N., Rasini E., Marino F. (2015). Influence of dopamine receptor gene polymorphisms on circulating T lymphocytes: A pilot study in healthy subjects. Human immunology, 76, 10, 747-752. <https://doi.org/10.1016/j.humimm.2015.09.032>

- Boydum A. Isolation of mononuclear cells and granulocytes from human blood. Scand.J.Clin.Lab. Invest. 21 (Suppl.97): 77-89, 1968

- Alex Mabou Tagne, Franca Marino, Massimiliano Legnaro, Alessandra Luini, Barbara Pacchetti and Marco Cosentino. A Novel Standardized Cannabis sativa L. Extract and Its Constituent Cannabidiol Inhibit Human Polymorphonuclear Leukocyte Functions. Int J Mol Sci 2019 Apr; 20(8): 1833. Published online 2019 Apr 13. doi: 10.3390/ijms20081833.

- Angela Scanzano, Laura Schembri, Emanuela Rasini, Alessandra Luini, Jessica Dallatorre, Massimiliano Legnaro, Raffaella Bombelli, Terenzio Congiu, Marco Cosentino, Franca Marino. Adrenergic Modulation of Migration, CD11b and CD18 Expression, ROS and interleukin-8 Production by Human Polymorphonuclear Leukocytes. Inflamm Res. 2015 Feb;64(2):127-35. doi: 10.1007/s00011-014-0791-8. Epub 2015 Jan 6.

DOI

dx.doi.org/10.17504/protocols.io.bpxsmpne

PROTOCOL CITATION

Marco Cosentino, Elisa Storelli, Alessandra Luini, Massimiliano LM Legnaro, Emanuela Rasini, Marco Ferrari, Franca Marino 2020. CELL COUNT- 02 - Manual cell count with Türk Solution. **protocols.io**
<https://dx.doi.org/10.17504/protocols.io.bpxsmpne>

LICENSE

———— This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

CREATED

Nov 21, 2020

LAST MODIFIED

Nov 21, 2020

PROTOCOL INTEGER ID

44754

MATERIALS TEXT

Istruments needed:

- Bürker chamber
- Optical microscope
- Plastic labware

Materials:

Gentian violet: Marco Viti

Acetic acid catalog number: A6283 Sigma Italy

ABSTRACT

Published work using this protocol:

- Kustrimovic N., Comi C., Magistrelli L., Rasini E., Legnaro M., Bombelli R., Aleksic I., Blandini F., Minafra B., Riboldazzi G., Struchio A., Mauri M., Bono G., Marino F., Cosentino M. Parkinson's disease patients have a complex phenotypic and functional Th1 bias: cross-sectional studies of CD4+ Th1/Th2/T17 and Treg in drug-naïve and drug-treated patients (2018). Journal of neuroinflammation, 15(1), 205. <https://doi.org/10.1186/s12974-018-1248-8>

- Kustrimovic, N., Rasini, E., Legnaro, M., Bombelli, R., Aleksic, I., Blandini, F., Comi, C., Mauri, M., Minafra, B., Riboldazzi, G., Sanchez-Guajardo, V., Marino, F., & Cosentino, M. (2016). Dopaminergic Receptors on CD4+ T Naive and Memory Lymphocytes Correlate with Motor Impairment in Patients with Parkinson's Disease. Scientific reports, 6, 33738. <https://doi.org/10.1038/srep33738>


- Cosentino M., Ferrari M., Kustrimovic N., Rasini E., Marino F. (2015). Influence of dopamine receptor gene polymorphisms on circulating T lymphocytes: A pilot study in healthy subjects. Human immunology, 76, 10, 747-752. <https://doi.org/10.1016/j.humimm.2015.09.032>

- Boydum A. Isolation of mononuclear cells and granulocytes from human blood. Scand.J.Clin.Lab. Invest. 21 (Suppl.97): 77-89, 1968

- Alex Mabou Tagne, Franca Marino, Massimiliano Legnaro, Alessandra Luini, Barbara Pacchetti and Marco Cosentino. A Novel Standardized Cannabis sativa L. Extract and Its Constituent Cannabidiol Inhibit Human Polymorphonuclear Leukocyte Functions. Int J Mol Sci 2019 Apr; 20(8): 1833. Published online 2019 Apr 13. doi: 10.3390/ijms20081833.


- Angela Scanzano, Laura Schembri, Emanuela Rasini, Alessandra Luini, Jessica Dallatorre, Massimiliano Legnaro, Raffaella Bombelli, Terenzio Congiu, Marco Cosentino, Franca Marino. Adrenergic Modulation of Migration, CD11b

- 1 Use Türk solution for checking cell purity.

Mix  **10 µl** of cell suspension with an equal amount of Türk solution (dilution factor = 2), allow mixture 3 min at room temperature.



SOLUTION- 08 - Türk solution
by **Farmacologia Medica**

- 2 Take  **10 µl** of the mixture and place it inside a Bürker chamber and view under an optical microscope using 40X magnification.
- 3 Count the cells in each square found in the four corners and in the central square (see figure 1 below), including those that lie on the bottom and left-hand perimeters, but not those that lie on the top and right hand perimeters (see figure 2 below).

Total number of cells per ml = mean number of cells x dilution factor x 10^4 (hemacytometer volume).




Figure 1

The gridded area of the chamber consists of nine 1 mmq squares. These squares are subdivided in three directions; 0.0625 mmq, 0.05 mmq and 0.04 mmq. The central square here in Figure 1 is further subdivided into 0.0025 mmq = 1/25 mmq squares. Count cells in 5 squares as shown.




Figure 2

Concerning those cells that lay on the perimeter of the square, count following this scheme.