

Sep 29, 2025

Microglia validation by FACS-analysis

DOI

dx.doi.org/10.17504/protocols.io.81wgbxokqlpk/v1

Riana Lo Bu^{1,2}, Frank Soldner^{1,2}

¹Albert Einstein College of Medicine, 1301 Morris Park Ave., Bronx, NY 10461, USA.;

²Aligning Science Across Parkinson's (ASAP) Collaborative Research Network, Chevy Chase, MD, 20815.



Oriol Busquets

Albert Einstein College of Medicine

OPEN ACCESS



DOI: https://dx.doi.org/10.17504/protocols.io.81wgbxokqlpk/v1

Protocol Citation: Riana Lo Bu, Frank Soldner 2025. Microglia validation by FACS-analysis. **protocols.io**

https://dx.doi.org/10.17504/protocols.io.81wgbxokqlpk/v1

License: This is an open access protocol distributed under the terms of the **Creative Commons Attribution License**, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited

Protocol status: Working

We use this protocol and it's working

Created: January 09, 2024

Last Modified: September 29, 2025

Protocol Integer ID: 93109

Keywords: ASAPCRN, microglia validation by fac, microglia validation, microglia validation by fluorescence, microglia differentiation protocol, procedure for microglial precursor, microglia differentiation protocol for cell generation, microglial precursor, microglia differentiation, activated cell sorting, cell sorting, fac, list of reagent

Funders Acknowledgements:

Aligning Science Across Parkinson's (ASAP)

Grant ID: ASAP-000486

Aligning Science Across Parkinson's (ASAP)

Grant ID: ASAP-024409



Abstract

This protocol describes the procedure for microglial precursor and mature microglia validation by Fluorescence-Activated Cell Sorting (FACS) analysis. Please refer to (https://doi.org/10.17504/protocols.io.4r3l22zbjl1y/v1) for microglia differentiation protocol for cell generation and media composition.

Protocol Overview

- A. Microglial precursor validation
- B. Mature microglia validation

General Notes

A list of reagents and relevant vendor information can be found in the table listed under the materials tab.

Attachments



Microglia validation...

41KB

Materials

Reagent Table:

| Item | Vendor | Catalog Number |
|-----------------------|------------|----------------|
| 1xPBS | Corning | MT21031CV |
| Accutase | Gemini | 400-158 |
| EDTA | Fisher | BP120500 |
| Trypan blue | Gibco | 15250061 |
| Human TruStain FcX | Biolegend | 422302 |
| APC Anti-human CD16 | Biolegend | 302102 |
| APC Anti-human CD45 | Biolegend | 304012 |
| APC anti-human CX3CR1 | Biolegend | 341610 |
| PE Anti-human P2ry12 | Biolegend | 392103 |
| PE Anti-human CD11b | Biolegend | 301306 |
| PE Anti-human CD14 | Fisher | 12014942 |
| Calcein violet | Invitrogen | C34858 |



Microglial precursor validation

52m

1 Harvest microglial precursors from flasks in STEP4 by collecting the media into 15 ml conical tubes and centrifuging it at \$\mathbb{\omega}\$ 150 rcf for \$\mathbb{\omega}\$ 00:04:00 .

4m

- 2 Remove the supernatant and resuspend the cells in STEP4 media.
- 3 Count the cells and aliquot 100,000 cells/experiment condition in 15ml conical tubes.

4m

- Remove the supernatant and resuspend the cells in Δ 100 μ L FACS buffer (PBS1x, EDTA 5 mM, 0.5% FBS).

10m

- Add primary antibodies for the desired experimental conditions ($\perp 1 \mu L$ /tube for the antibodies described above or according to manufacturer's recommendations).
- 8 Incubate for 500:30:00 on 600 ice.

30m

9 Wash the cells 2 times by adding 4 2 mL /tube of FACS buffer followed by centrifugation at 150 rcf for 00:04:00

4m

- 10 Resuspend the cells in Δ 350 μ L of FACS buffer + calcein violet (follow manufacturer's instructions as to how to prepare calcein violet).
- 11 Keep your samples protected from light on ice and perform standart FACS analysis.



Mature Microglia validation

32m

- Remove the media from microglia cultures and add accutase (4 1 mL /well in 6 well plates).
- 13 Incubate (5) 00:10:00 at \$ 37 °C.

10m

- Add <u>Add</u> 3 mL of microglia maturation media to quench the reaction and collect the cells in 15 ml conical tubes.
- 15 Spin down at \$\mathbb{3}\$ 150 rcf for \mathbb{\infty} 00:08:00 \ .

8m

- Remove the supernatant and resuspend the cells in microglia maturation media.
- 17 Count the cells and aliquot 100,000 cells/experiment condition in 15ml conical tubes.

4m

- Remove the supernatant and resuspend the cells in \perp 100 μ L FACS buffer (PBS1x, EDTA 5 mM, 0.5% FBS).
- Add Δ 5 μ L /tube of human TruStain FcX and incubate for 00:10:00 at Room temperature .

10m

- Add primary antibodies for the desired experimental conditions ($\Delta 1 \mu L$ /tube for the antibodies described above or according to manufacturer's recommendations).

30m



- 23 Wash the cells 2 times by adding with by adding ___ 2 mL /tube of FACS buffer followed
- 4m
- 24 Resuspend the cells in \perp 350 μ L of FACS buffer + calcein violet (follow manufacturer's instructions as to how to prepare calcein violet).
- 25 Keep your samples protected from light on in ice and perform standard FACS analysis.

Protocol references

https://doi.org/10.17504/protocols.io.4r3l22zbjl1y/v1