



NOV 13, 2023

OPEN ACCESS



**DOI:**  
[dx.doi.org/10.17504/protocol.s.io.81wgbxqkolpk/v1](https://dx.doi.org/10.17504/protocol.s.io.81wgbxqkolpk/v1)

**External link:**  
<https://docs.google.com/document/d/1bvNLoEJIWpkJ6-heG1qyHc5Z6WEqOAqlsZwootmQpm8/edit>

**Protocol Citation:** Shlomo Yakir Hoch, Ravit Netzer, Karen Hakeny, Sarel J Fleishman 2023. GGAssembler Library Construction. **protocols.io** <https://dx.doi.org/10.17504/protocols.io.81wgbxqkolpk/v1>

**License:** This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited


**Created:** Nov 12, 2023

## GGAssembler Library Construction

Forked from [PCR with Q5® Polymerase \(M0491\)](#)

Shlomo Yakir Hoch<sup>1</sup>, Ravit Netzer<sup>1,2</sup>, Karen Hakeny<sup>1</sup>, Sarel J Fleishman<sup>1,2</sup>

<sup>1</sup>Weizmann Institute of Science; <sup>2</sup>Scala Biodesign LTD



Shlomo Yakir Hoch

### ABSTRACT

This protocol describes methods for GGAssembler.

### GUIDELINES

**Please note that protocols with Q5® High-Fidelity DNA Polymerase may differ from protocols with other polymerases. Conditions recommended below should be used for optimal performance.**

#### Reaction Setup:

We recommend assembling all reaction components on ice and quickly transferring the reactions to a thermocycler preheated to the denaturation temperature (98°C). All components should be mixed prior to use. Q5 High-Fidelity DNA Polymerase may be diluted in 1X Q5 Reaction Buffer just prior to use in order to reduce pipetting errors.

#### General Guidelines:

##### 1. Template:

Use of high quality, purified DNA templates greatly enhances the success of PCR. Recommended amounts of DNA template for a 50 µl reaction are as follows:

A	B
DNA Genomic	1 ng–1 µg
Plasmid or Viral	1 pg–1 ng

##### 2. Primers:

Oligonucleotide primers are generally 20–40 nucleotides in length and ideally have a GC content of 40–60%. Computer programs such as [Primer3](#) can be used to design or analyze primers. The best results are typically seen when using each primer at a

**Last Modified:** Nov 13, 2023

**PROTOCOL integer ID:** 90809

**Funders**

**Acknowledgement:**

Volkswagen Foundation  
Grant ID: 94747

Israel Science Foundation  
Grant ID: 1844

European Research Council  
through a Consolidator Award  
Grant ID: 815379

Dr. Barry Sherman Institute  
for Medicinal Chemistry  
Donation in memory of Sam  
Switzer

final concentration of 0.5  $\mu$ M in the reaction.

3.  $Mg^{++}$  and additives:

$Mg^{++}$  concentration of 2.0 mM is optimal for most PCR products generated with Q5 High-Fidelity DNA Polymerase. When used at a final concentration of 1X, the Q5 Reaction Buffer provides the optimal  $Mg^{++}$  concentration.

Amplification of some difficult targets, like GC-rich sequences, may be improved by the addition of 1X Q5 High GC Enhancer. The Q5 High GC Enhancer is not a buffer and should not be used alone. It should be added only to reactions with the Q5 Reaction Buffer when other conditions have failed.

4. Deoxynucleotides:

The final concentration of dNTPs is typically 200  $\mu$ M of each deoxynucleotide. Q5 High-Fidelity DNA Polymerase cannot incorporate dUTP and is not recommended for use with uracil-containing primers or templates.

5. Q5 High-Fidelity DNA Polymerase concentration:

We generally recommend using Q5 High-Fidelity DNA Polymerase at a final concentration of 20 units/ml (1.0 unit/50  $\mu$ l reaction). However, the optimal concentration of Q5 High-Fidelity DNA Polymerase may vary from 10–40 units/ml (0.5–2 units/50  $\mu$ l reaction) depending on amplicon length and difficulty. Do not exceed 2 units/50  $\mu$ l reaction, especially for amplicons longer than 5 kb.

6. Buffers:

The 5X Q5 Reaction Buffer provided with the enzyme is recommended as the first-choice buffer for robust, high-fidelity amplification. For difficult amplicons, such as GC-rich templates or those with secondary structure, the addition of the Q5 High GC Enhancer can improve reaction performance. The 5X Q5 Reaction Buffer is detergent-free and contains 2.0 mM  $Mg^{++}$  at the final (1X) concentration.

7. Denaturation:

An initial denaturation of 30 seconds at 98°C is sufficient for most amplicons from pure DNA templates. Longer denaturation times can be used (up to 3 minutes) for templates that require it.

During thermocycling, the denaturation step should be kept to a minimum. Typically, a 5–10 second denaturation at 98°C is recommended for most templates.

8. Annealing:

Optimal annealing temperatures for Q5 High-Fidelity DNA Polymerase tend to be higher than for other PCR polymerases. The [NEB  \$T\_m\$  Calculator](#) should be used to determine the annealing temperature when using this enzyme. Typically, use a 10–

30 second annealing step at 3°C above the  $T_m$  of the lower  $T_m$  primer. A temperature gradient can also be used to optimize the annealing temperature for each primer pair.

For high  $T_m$  primer pairs, two-step cycling without a separate annealing step can be used (see note 11).

#### 9. Extension:

The recommended extension temperature is 72°C. Extension times are generally 20–30 seconds per kb for complex, genomic samples, but can be reduced to 10 seconds per kb for simple templates (plasmid, *E. coli*, etc.) or complex templates < 1 kb. Extension time can be increased to 40 seconds per kb for cDNA or long, complex templates, if necessary.

A final extension of 2 minutes at 72°C is recommended.

#### 10. Cycle number:

Generally, 25–35 cycles yield sufficient product. For genomic amplicons, 30–35 cycles are recommended.

#### 11. 2-step PCR:

When primers with annealing temperatures  $\geq 72^\circ\text{C}$  are used, a 2-step thermocycling protocol (combining annealing and extension into one step) is possible.

#### 12. Amplification of long products:

When amplifying products > 6 kb, it is often helpful to increase the extension time to 40–50 seconds/kb.

#### 13. PCR product:

The PCR products generated using Q5 High-Fidelity DNA Polymerase have blunt ends. If cloning is the next step, then blunt-end cloning is recommended. If T/A-cloning is preferred, the DNA should be purified prior to A-addition, as Q5 High-Fidelity DNA Polymerase will degrade any overhangs generated.







Addition of an untemplated -dA can be done with *Taq* DNA Polymerase ([NEB #M0267](#)) or Klenow  $\text{exo}^-$  ([NEB #M0212](#)).

## MATERIALS

### Materials



Q5 High-Fidelity DNA Polymerase - 100 units New England  
Biolabs Catalog #M0491S


-  DpnI - 5,000 units New England Biolabs Catalog #R0176L
-  T4 DNA Ligase Reaction Buffer - 6.0 ml New England Biolabs Catalog #B0202S
-  T4 DNA Ligase - 20,000 units New England Biolabs Catalog #M0202S
-  BsaI-HFv2 New England Biolabs Catalog # R3733S
-  Nuclease-free Water - 25 ml New England Biolabs Catalog #B1500S
-  Monarch® PCR & DNA Cleanup Kit (5 µg) New England Biolabs Catalog #T1030

Equipment

Equipment	
	NAME
Qubit 2.0 Fluorometer instrument	BRAND
Q33226	SKU
with Qubit RNA HS Assays	SPECIFICATIONS

Equipment	
4200 TapeStation System	NAME
Electrophoresis tool for DNA and RNA sample quality control.	TYPE
TapeStation Instruments	BRAND
G2991AA	SKU
<a href="https://www.agilent.com/en/product/automated-electrophoresis/tapestation-systems/tapestation-instruments/4200-tapestation-system-228263">https://www.agilent.com/en/product/automated-electrophoresis/tapestation-systems/tapestation-instruments/4200-tapestation-system-228263</a>	LINK

## SAFETY WARNINGS

 Please refer to the Safety Data Sheets (SDS) for health and environmental hazards.

## BEFORE START INSTRUCTIONS

Please note that protocols with Q5® High-Fidelity DNA Polymerase may differ from protocols with other polymerases. Conditions recommended below should be used for optimal performance.

### Amplify constant fragments

**1** Set up the following reaction  On ice :


#### Note

All components should be mixed prior to use. Q5 High-Fidelity DNA Polymerase may be diluted in 1X Q5 Reaction Buffer just prior to use in order to reduce pipetting errors.

A	B	C	D
COMPONENT	25 µl REACTION	50 µl REACTION	FINAL CONCENTRATION
5X Q5 Reaction Buffer	5 µl	10 µl	1X
10 mM dNTPs	0.5 µl	1 µl	200 µM
10 µM Forward Primer	1.25 µl	2.5 µl	0.5 µM
10 µM Reverse Primer	1.25 µl	2.5 µl	0.5 µM
Template DNA	variable	variable	< 1,000 ng
Q5 High-Fidelity DNA Polymerase	0.25 µl	0.5 µl	0.02 U/µl
5X Q5 High GC Enhancer (optional)	(5 µl)	(10 µl)	(1X)
Nuclease-Free Water	to 25 µl	to 50 µl	

**2** Gently mix the reaction.

3 Collect all liquid to the bottom of the tube by a quick spin if necessary and overlay the sample with mineral oil if using a PCR machine without a heated lid.

4 Quickly transfer PCR tubes to a PCR machine preheated to the denaturation temperature (  98 °C ) and begin thermocycling.



**Thermocycling Conditions for a Routine PCR:**

A	B	C
STEP	TEMP	TIME
Initial Denaturation	98°C	30 seconds
25–35 Cycles	98°C	5–10 seconds
	*50–72°C	10–30 seconds
	72°C	20–30 seconds/kb
Final Extension	72°C	2 minutes
Hold	4–10°C	

\*Use of the [NEB Tm Calculator](#) is highly recommended.

5 (IF REQUIRED) Digest the lead gene templates with DpnI.



6 Quantify DNA concentration by Qubit.

7 Measure DNA length using TapeStation.

## Variable fragments fill in

- 8 Follow the steps for Amplify constant fragments (steps 1-7).
- 9 Combine variable fragments of the same segment in equal concentration.



## Golden Gate Assembly

- 10 Set up 25 µl assembly reactions as follows:



REAGENTS	ASSEMBLY REACTION
DNA inserts 100 ng/ul each	1 µl (~ 100 ng) each, (up tp 19.5 µl)
T4 DNA Ligase Buffer (NEB #B0202) (10X)	2.5 µl
T4 DNA Ligase (NEB #M0202), 2000 U/µl	0.5 µl (1000 units)
BsaI-HFv2 (NEB #R3733), 20 U/µl	1.5 µl (30 units)
Nuclease-free H2O (NEB #B1500)	to 25 µl

- 11 Mix gently by pipetting up and down 4 times.



- 12 Briefly microcentrifuge (1 second) to bring material to the bottom of tube.



- 13 Transfer to thermocycler and program as follows: (5 min 37°C → 5 min 16°C) x 30 cycles followed by 5 minutes 60°C. If reactions are done overnight, add a 4°C terminal hold to the protocol, but repeat the final 5 minutes 60°C step the next day before the transformations.



## Golden Gate Assembly Cleanup

14



### Note

This step is optional but HIGHLY recommended.

Perform Oligonucleotide Cleanup (<https://dx.doi.org/10.17504/protocols.io.bg9sjz6e>) or follow a size selection protocol.

## Amplify Golden Gate Assembly Products

15



For 1x reaction set up the following reaction  On ice :

### Note

Using GG assembly products in concentration of 3-5ng/ $\mu$ l is highly recommended.

A	B	C
Component	25 $\mu$ l Reaction	Final Concentration
Q5 High-Fidelity 2X Master Mix	12.5 $\mu$ l	1X
10 $\mu$ M Forward Primer	1.25 $\mu$ l	0.5 $\mu$ M
10 $\mu$ M Reverse Primer	1.25 $\mu$ l	0.5 $\mu$ M
GG assembly products	5 $\mu$ l	
Nuclease-Free Water	to 25 $\mu$ l	

16



Gently mix the reaction.




17



Collect all liquid to the bottom of the tube by a quick spin if necessary and overlay the sample with mineral oil if using a PCR machine without a heated lid.

18



Quickly transfer PCR tubes to a PCR machine preheated to the denaturation temperature (  98 °C ) and begin thermocycling.

#### Thermocycling Conditions for a Routine PCR:

A	B	C
STEP	TEMP	TIME
Initial Denaturation	98°C	30 seconds
25–35 Cycles	98°C	5–10 seconds
	*50–72°C	10–30 seconds
	72°C	20–30 seconds/kb
Final Extension	72°C	2 minutes
Hold	4–10°C	

\*Use of the [NEB Tm Calculator](#) is highly recommended.

## Golden Gate Assembly Amplification Cleanup

19



#### Note


This step is optional but HIGHLY recommended.

Perform Oligonucleotide Cleanup (<https://dx.doi.org/10.17504/protocols.io.bg9sjz6e>) or follow a size selection protocol.

## Golden Gate Assembly Amplification Quantification

20

Quantify DNA concentration by Qubit.



## 21 Measure DNA length using TapeStation.