



Apr 17, 2020

Simple Step to Increase Gibson Assembly Efficiency ^{v.1}

Eric Danner¹¹Max Delbrück Center
1 Works for me dx.doi.org/10.17504/protocols.io.7kzhkx6

Eric Danner
 Max Delbrück Center


ABSTRACT

The Gibson assembly solution is extremely toxic to some bacteria cells.

Doing a solution change after assembly and before transforming the bacteria will significantly improve bacteria survival. This increases 10-100x the amount of colonies.

MATERIALS

NAME	CATALOG #	VENDOR
Gibson Assembly Master Mix - 50 rxns	E2611L	New England Biolabs
NucleoSpin® Gel and PCR Clean-up	740609.10	Macherey and Nagel

Do Normal Assembly

- 1 Do a normal Gibson assembly. Incubate in the PCR machine 60 min @ 50C
- 2 Remove the assemblies. Do a PCR quickchange/ gel extract column. This will remove the Gibson Assembly solution but keep the 100ng (or however much) of DNA.

Elute in 15u -20ul TE or water
- 3 Take the 15-20ul assembled plasmid and heat shock your bacteria. 30 second heat shock. 2 min ice. Add 200ul media and shake for 45 min. Carb can be plated immediately.

Plate



This is an open access protocol distributed under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited